



WEBINAR HOUSEKEEPING

Attendee Participation

Open your control panel.


Join audio:

- Choose **Phone Call** and dial using the information provided.
- Or choose **Computer Audio** to use your computer's speakers for audio.
- Attendee audio will be muted.

Submit questions and comments via the Questions panel. If you would like to unmute and ask your questions, please specify that in the Questions Panel.

If viewing this webinar as a group, please provide a list of attendees via the Questions panel.

Note: Today's presentation is being recorded and will be provided on the FTP after the webinar.





CRYSTAL RIVER/KINGS BAY, HOMOSASSA/CHASSAHOWITZKA SPRINGS, AND WEEKI WACHEE BASIN MANAGEMENT ACTION PLAN UPDATES

Chandler Keenan, Environmental Consultant
Sam Hankinson, Professional Geologist II
Water Quality Restoration Program
Florida Department of Environmental Protection

GoTo Webinar | May 30, 2024



AGENDA

- Basin Management Action Plan (BMAP) Overview.
- Nitrogen Source Inventory Loading Tool (NSILT):
 - Updates.
 - Draft Results.
- Spring Vent Load Analysis.
- Next Steps - BMAP Updates:
 - Draft Allocation Approach.
 - Milestones.






BMAPs



One of DEP's methods for **restoring water quality** in an impaired waterbody.

BMAPs are:

- Developed with stakeholder input.
- Adopted by The Florida Department of Environmental Protection's (DEP) Secretarial Order.
- Enforceable.
- Implemented through a phased approach.
- Reported on annually.
- Updated regularly.





KEY BMAP COMPONENTS

- Total maximum daily loads (TMDLs) being addressed.
- Area addressed by the restoration plan.
- Identify sources.
- Phased implementation approach.
- Milestones.
- Projects and management strategies.
- Future growth impacts.

Projects to meet the TMDL:

- Implementation timeline.
- Commitment to projects.
- Expected water quality improvement from projects and management strategies.




Process to assess progress toward achieving the TMDL:

- Monitoring plan.
- Project reporting.
- Periodic follow-up meetings.
- Water quality analyses.



BMAPs

SPRINGS COAST OVERVIEW



Background

Early 2000s - Many of Florida's freshwater springs determined to be impaired for nutrients.

2014 - DEP adopted TMDLs for many waterbodies in Springs Coast basin.

2016 - Florida Legislature designated 30 Outstanding Florida Springs (OFS) to require additional protections.

2018 - BMAPs adopted as restoration framework to meet TMDLs.


Today - Working on updates to the Springs BMAPs to be adopted by 2025.



CRYSTAL RIVER / KINGS BAY BMAP

- BMAP area is approximately 178,753 acres and includes OFS including Kings Bay Group (Hunter, House, Black, Idiot's Delight, and Tarpon Hole Springs).
- Impaired for the nitrate form of nitrogen and orthophosphate.
- Total maximum daily loads (TMDLs) are
 - Springs 0.23 mg/L nitrate annual average.
 - Kings Bay 0.28 mg/L nitrate annual average.
 - Springs 0.028 mg/L orthophosphate annual average.
 - Kings Bay 0.032 mg/L orthophosphate annual average.

Type of Entity	Name
Responsible Stakeholders	Citrus County City of Crystal River Agricultural producers Golf courses
Responsible Agencies	Florida Department of Agriculture and Consumer Services Florida Department of Environmental Protection Florida Department of Health Southwest Florida Water Management District
Other Interested Stakeholders	Citizens Duke Energy Florida Farm Bureau Federation Florida Onsite Wastewater Association Gulf Archeology Research Institute Homeowners/Citizens Kings Bay Rotary Kings Bay Springs Alliance Save Crystal River Save the Manatee Club St. Martins Marsh Aquatic Preserve University of Florida Institute of Food and Agricultural Sciences – Citrus County Extension Service U.S. Fish and Wildlife Service






HOMOSASSA/ CHASSAOWITZKA SPRINGS BMAP

- BMAP area is approximately 340,609 acres and include OFS Homosassa and Chassahowitzka Springs Groups.
- Impaired for the nitrate form of nitrogen.
- Total maximum daily loads (TMDLs) are nitrate annual averages of 0.23 mg/L for the springs and 0.25 mg/L for the rivers.

Type of Entity	Name
Responsible Stakeholders	Citrus County City of Brooksville City of Inverness Hernando County Agricultural producers Golf courses
Responsible Agencies	Florida Department of Agriculture and Consumer Services Florida Department of Environmental Protection Florida Department of Health Southwest Florida Water Management District
Other Interested Stakeholders	Citizens/Homeowners Florida Farm Bureau Florida Onsite Wastewater Association Florida Springs Council Hernando Beach Government Affairs Committee Hernando County Port Authority Hernando Environmental Land Protectors (HELP) Homosassa River Alliance Save the Manatee Club






WEEKI WACHEE BMAP

- BMAP area is approximately 200,474 acres and includes the Weeki Wachee OFS.
- Impaired for the nitrate form of nitrogen.
- Total maximum daily loads (TMDLs) are nitrate annual averages of 0.28 mg/L for Weeki Wachee Spring, 0.23 mg/L for the Magnolia-Aripeka Spring Groups, and 0.20 mg/L for the Weeki Wachee River.

Type of Entity	Name
Responsible Stakeholders	City of Brooksville Hernando County Pasco County Agricultural producers Golf courses
Responsible Agencies	Florida Department of Agriculture and Consumer Services Florida Department of Environmental Protection Florida Department of Health Southwest Florida Water Management District
Other Interested Stakeholders	Citizens City of Weeki Wachee Florida Farm Bureau Florida Onsite Wastewater Association Florida Springs Institute Hernando Beach Government Affairs Committee Hernando County Task Force Hernando Environmental Land Protectors (HELP) Save the Manatee Club





CLEAN WATERWAYS ACT: TIMELINE

June 12, 2023

Final Order signed by the Secretary.



July 12, 2023

Deadline for written explanation of potential exemption to be submitted to the department.



Feb. 1, 2024

Deadline for submitting draft Onsite sewage treatment and disposal system (OSTDS) remediation and/or wastewater treatment plans for the department's review.



Aug. 1, 2024

Deadline for submitting complete OSTDS remediation and/or wastewater treatment plans to the department.

The nutrient BMAPs included in the Final Order require these plans.



HOUSE BILL (HB) 1379: ENVIRONMENTAL PROTECTION

Increased protection for OFS.


Strengthens Water Quality Protections and BMAPs.

HB 1379

Improves Local Government Long-Term Comprehensive Planning.

Expands Funding Opportunities to Address Water Quality Impairments.

Expanded prohibitions in OFS to entire BMAP area. (section 373.811, Florida Statutes [F.S.]



- New conventional OSTDS where sewer is available.
- New domestic wastewater disposal facilities with permitted capacities of 100,000 gallons per day or more, unless they meet Advanced Waste Treatment (AWT) standards.
- New HAZMAT disposal facilities.
- Land application of Class A or B biosolids not in accordance with a DEP-approved nutrient management plan.
- New agricultural operations not implementing best management practices (BMPs), measures necessary to achieve pollution reduction levels or groundwater monitoring plans.



2024 DEP AGENCY BILL: HB 1557

Advances the protection of our environmental resources by:

- Enhancing protections for Florida's Coral Reef and aquatic preserves.
- Strengthening Florida's resilience planning.
- Furthering the Onsite Septic Program transfer.

Improving Treatment of Reclaimed Water

Ensures that reclaimed water is treated to meet AWT or a more stringent treatment standard in certain BMAP areas, while still promoting its use to eliminate surface water discharges and meet water supply challenges.

Expanding Wastewater Facility Plans

Supports the development of domestic wastewater treatment plans and OSTDS remediation plans within BMAP or other restoration areas by requiring facilities to provide information to the local entities developing these plans.

Investing in Innovative Technologies

Creates a program to expeditiously review new and innovative enhanced nutrient-reducing OSTDS to reduce the nutrients entering Florida's waterways.




STAR

STATEWIDE ANNUAL REPORT – PROJECT REPORTING

What is the STAR?

- Summarizes accomplishments in the BMAPs statewide.
- Reports on restoration projects and management strategies.
- Published July 1 of each year.
- Currently in the process of project updates and verification for STAR 2023.





PRELIMINARY STAR RESULTS FOR 2023


BMAP	Project Count *					Verified Project Reductions of Total Nitrogen
	Planned	Ongoing	Underway	Completed	Total	
Kings Bay/ Crystal River	20	7	10	34	71	23,770 lbs./yr.
Homosassa/ Chassahowitzka	12	12	19	25	68	19,056 lbs./yr.
Weeki Wachee	18	16	11	36	81	61,083 lbs./yr.

*As of Dec. 31, 2023



WATER QUALITY DATA




KINGS BAY NITRATE DATA





WATER QUALITY DATA

HOMOSASSA/CHASSAHOWITZKA AND WEEKI WACHEE DATA





DATA UPLOAD

WATERSHED INFORMATION NETWORK (WIN)

- Through both the WIN and Florida STORET (STOrage and RETrieval) data repositories, DEP implements Florida statutory requirements, DEP rule requirements and U.S. Environmental Protection Agency (EPA) funding requirements for management of environmental (non-regulatory) data for the state.
- Data from WIN are used by DEP for standards development, Impaired Waters Rule assessments, TMDL development, reasonable assurance plans, alternative restoration plans, **BMAP development and assessment** and for providing data as required to EPA and to the public.
- WIN data can be retrieved through the WIN Reports and Extracts menu at <https://prodenv.dep.state.fl.us/DearWin/>.
- Data providers to WIN and STORET include Division of Environmental Assessment and Restoration and other DEP entities, water management districts (WMDs), cities, counties, other state agencies, universities, private and volunteer organizations.
- If your entity is collecting ambient water quality data, please upload it to WIN.



WIN COORDINATORS


WIN Coordinator	DEP District Area or Role	Phone	Email
Justin Nelson	Northeast, Northwest, Southeast	850-245-8510	Justin.M.Nelson@FloridaDEP.gov
Casey Marston	South, Southwest	850-245-8049	Casey.Marston@FloridaDEP.gov
Lisa Schwenning	SPA (STORET Public Access), WQX (U.S. EPA Water Quality Exchange)	850-245-8509	Lisa.Schwenning@floridaDEP.gov
Jason Storrs	Central, Statewide	850-245-8467	Jason.Storrs@FloridaDEP.gov



BMAP UPDATES

ADOPTED BY JULY 1, 2025

- NSILT updates.
- Spring vent load analyses.
- Entity allocation development.
- Future growth.
- Establish five-year milestones for project implementation.
- Incorporate additional projects.
- Incorporate Clean Waterways Act (Senate Bill 712) requirements.
- Incorporate HB 1379 requirements.
- Incorporate regional projects.
- Water quality data evaluation:
 - Evaluation of the monitoring network (spring vent and groundwater).
 - Water quality trend analyses.
- Evaluate further OSTDS provisions.
- Evaluate AWT or other more stringent effluent limits.
- Update the BMAP documents.






BMAP MEETING

PUBLIC QUESTIONS PERIOD

Verbal Questions:

- We ask that questions and comments be limited to **3 minutes** so that we may hear from everyone.
- Please type in the chat if you would like to unmute yourself and ask a question or comment.



Written Comments:

- Submit written comments concerning today's meeting to BMAPProgram@FloridaDEP.gov.





NSILT UPDATES

- NSILT Process.
- Methodology review for sources.
 - Atmospheric deposition (AD).
 - Wastewater treatment facilities (WWTFs).
 - OSTDS.
 - Urban turfgrass fertilizer (UTF).
 - Sports turfgrass fertilizer (STF).
 - Farm fertilizer (FF).
 - Livestock waste (LW).
 - Biosolids.
- Draft results.





NSILT GENERAL PROCESS SUMMARY

Estimate loading to land surface for each source category.


Apply a source specific, literature derived biochemical attenuation factor to surface loading estimate.

Apply a location specific recharge factor to surface loading estimate.

LOADING TO GROUNDWATER.



NITROGEN CYCLE AND ATTENUATION





RECHARGE TO GROUNDWATER


- Divided into four recharge categories based on a composite recharge map of the Floridan aquifer.
- The recharge amount evaluates the percent of nitrogen loading that is expected to impact spring vent after bioattenuation is considered. Recharge rates are summarized in the table below for each recharge category.
- Delineation of recharge areas and associated recharge rates are consistent with the previous NSILT report.
- Recharge factors are applied to estimate loading for all source categories based on location of deposition.

Recharge to Groundwater	
Category	Rate
High	90%
Medium	50%
Low	10%
Discharge	0%




RECHARGE TO GROUNDWATER


Kings Bay/Crystal River



Homosassa/Chassahowitzka Springs



Weeki Wachee





AD


- Estimated using a nationwide model developed by the Total Deposition Science Committee and EPA called the Total Deposition (TDEP) model.
- AD estimates from 2019 and 2020 were averaged to estimate annual loading in the springsheds.
- Methodology is consistent with previous NSILT.

Springshed	Estimated Load to Surface (lbs-N/yr)	Estimated Load to Groundwater (lbs-N/yr)
Kings Bay	814,302	69,099
Homosassa	920,811	70,808
Chassahowitzka	545,574	43,944
Weeki Wachee	1,147,400	93,069




AD


Kings Bay/Crystal River



Homosassa/Chassaowitzka Springs



Weeki Wachee





WWTFs


- Wastewater Facility Regulation information was reviewed to determine the location of all WWTFs, as well as their effluent application or disposal sites.
- Discharge monitoring report data collected by Wastewater Facility Regulation from 2019 to 2021 were reviewed to determine effluent discharge volume and total nitrogen (TN) concentration for each disposal stream.
- Effluent disposal sites evaluated include reuse, disposal in a rapid infiltration basin, sprayfield, absorption field or wetland. Each method has its own estimated attenuation factor.

Wastewater Treatment Effluent Attenuation				
Reuse	RIB	Sprayfield	Absorption Field	Wetland
75%	25%	60%	50%	85%



WWTFs

KINGS BAY/CRYSTAL RIVER




Kings Bay					
County	Facility ID	Facility Name	Disposal Type	Recharge Type at Disposal Site	Gross TN Average Load (lbs/yr)
Citrus	FLA011844	Brentwood Regional WWTF	RIB	High	9731.73
Citrus	FLA011845	Meadowcrest WWTF	RIB	High	10459.22
Citrus	FLA011845	Meadowcrest WWTF	Reuse (golf, resi)	High	10030.46
Citrus	FLA011846	New Horizons WWTF	RIB	High	196.42
Citrus	FLA011848	Crystal River City Of WWTF	Sprayfield	High	8508.59
Citrus	FLA011848	Crystal River City Of WWTF	Reuse (golf, resi)	High	7969.03
Citrus	FLA011849	Crystal Acres MHP WWTF	RIB	High	80.88
Citrus	FLA011849	Crystal Acres MHP WWTF	Sprayfield	High	40.67
Citrus	FLA011854	Pelican Bay Apartments	RIB	Medium	239.41
Citrus	FLA011855	Sandy Oaks RVP & MHC WWTF	Sprayfield	High	81.87
Citrus	FLA011855	Sandy Oaks RVP & MHC WWTF	RIB	High	81.87
Citrus	FLA011861	Bayfront Health Seven Rivers	RIB	Medium	554.05
Citrus	FLA011863	Lake Rousseau Resort LLC	RIB	High	72.35
Citrus	FLA011869	Beverly Hills WWTF	RIB	High	20508.95
Citrus	FLA011876	Indian Springs Utilities	RIB	Medium	850.13
Citrus	FLA011895	Thunderbird MHP WWTF	RIB	Medium	148.38
Citrus	FLA011914	Greenbriar Of Citrus Hills	RIB	High	517.51
Citrus	FLA011918	Citrus Center Shopping Center WWTF	RIB	High	443.91
Citrus	FLA011920	Inverness Park	Sprayfield	High	210.6
Citrus	FLA011922	Quality Inn	RIB	Medium	344.74
Citrus	FLA011924	Lecanto Hills MHP WWTF	RIB	High	83.03
Citrus	FLA011928	Ventura Village Apartments WWTF	Absorption Field	High	169.97
Citrus	FLA011872	Imperial Gardens MHP	RIB	Medium	73.42



WWTFs

HOMOSASSA/CHASSAHOWITZKA SPRINGS




Homosassa Spring Group					
County	Facility ID	Facility Name	Disposal Type	Recharge Type at Disposal Site	Gross TN Average Load (lbs/yr)
Citrus	FLA011839	Floral City Elementary School WWTF	RIB	High	44.48
Citrus	FLA011847	Inverness, City of - WWTF	Sprayfield	High	2,712.03
Citrus	FLA011847	Inverness, City of - WWTF	Reuse (golf, resi)	High	1,396.84
Citrus	FLA011847	Inverness, City of - WWTF	Reuse (golf, resi)	Medium	52.83
Citrus	FLA011853	Aunt Vera's Antique Store	RIB	Medium	4.23
Citrus	FLA011864	Moonrise Resort	RIB	High	222.5
Citrus	FLA011879	Oak Pond Mobile Home Estates	RIB	Low	126.99
Citrus	FLA011880	Stoneridge Landing	RIB	High	309.33
Citrus	FLA011884	Floral Oaks Apartments WWTF	RIB	High	51.08
Citrus	FLA011891	Bedrock Singing Forest MHP WWTF	RIB	High	469.56
Citrus	FLA011893	Point O Woods	RIB	High	346.83
Citrus	FLA011893	Point O Woods	Reuse (golf, resi)	High	567.4
Citrus	FLA011898	Harbor Lights Mobil Home Resort WWTP	Sprayfield	High	108.56
Citrus	FLA011899	Cedar Lakes MHP WWTF	RIB	Medium	396.41
Citrus	FLA011900	Royal Oaks Manor	RIB	High	423.83
Citrus	FLA011901	Bell Villa MHP	RIB	Discharge	144.47
Citrus	FLA011901	Bell Villa MHP	RIB	Low	30.42
Citrus	FLA011904	Oasis Mobile Home and RV Park WWTF	RIB	High	60.08
Citrus	FLA011907	Evanridge MHP	RIB	Low	167.41
Citrus	FLA011915	Forest View MHP WWTF	RIB	Medium	700.78
Citrus	FLA011927	Tarawood Adult Community	RIB	High	166.06
Citrus	FLA011902	Palm Terrace Village WWTF	Sprayfield	High	186.68
Chassahowitzka Spring Group Spring Group					
County	Facility ID	Facility Name	Disposal Type	Recharge Type at Disposal Site	Gross TN Average Load (lbs/yr)
Citrus	FLA011903	Sugarmill Woods WWTF	Sprayfield	High	10689.67
Citrus	FLA011903	Sugarmill Woods WWTF	Sprayfield	Medium	2815.03
Citrus	FLA011903	Sugarmill Woods WWTF	RIB	High	18177.11
Citrus	FLA011916	Walden Woods MHP WWTF	RIB	High	741.98
Hernando	FLA012071	Wesleyan Village	RIB	High	934.23
Hernando	FLA012071	Wesleyan Village	Sprayfield	High	49.25
Hernando	FLA012062	Countryside Estates WWTF	RIB	High	151.23
Hernando	FLA012046	Brooksville Golf & Country Club WWTF	RIB	High	58.6



WWTFs

WEEKI WACHEE



Weeki Wachee Spring Group					
County	Facility ID	Facility Name	Disposal Type	Recharge Type at Disposal Site	Gross TN Average Load (lbs/yr)
Hernando	FLA012036	William S. Smith Water Reclamation Facility	Reuse (golf, resi)	Medium	13731.60934
Hernando	FLA012036	William S. Smith Water Reclamation Facility	Reuse (golf, resi)	High	10130.48234
Hernando	FLA012038	Weeki Wachee North MHP WWTF	RIB	High	413.0892313
Hernando	FLA012039	Eckerd Acedemy at Brooksville WWTF	RIB	High	152.1139462
Hernando	FLA012043	Spring Hill WRF	RIB	Discharge	6861.160315
Hernando	FLA012043	Spring Hill WRF	Reuse (golf, resi)	High	52957.23045
Hernando	FLA012044	Camp - A - Wyle Resort WWTF	RIB	High	334.0131079
Hernando	FLA012045	Camper's Holiday Association WWTF	RIB	Medium	221.4558765
Hernando	FLA012045	Camper's Holiday Association WWTF	Absorption Field	Medium	222.2516069
Hernando	FLA012048	Big Tree MH & RV Village	Absorption Field	High	333.8881498
Hernando	FLA012048	Big Tree MH & RV Village	RIB	High	0
Hernando	FLA012054	Frontier Campground MHP	RIB	High	306.4342082
Hernando	FLA012059	Imperial Estates MHP WWTF	RIB	High	388.6142913
Hernando	FLA012059	Imperial Estates MHP WWTF	Sprayfield	High	365.3332508
Hernando	FLA012065	Topics RV Community WWTF	RIB	High	153.6426252
Hernando	FLA012066	Camp Sonshine WWTF (fka Church of God of Pro	RIB	High	71.53348156
Hernando	FLA012069	Glen Water Reclamation Facility	RIB	High	18971.44336
Hernando	FLA012069	Glen Water Reclamation Facility	Sprayfield	High	11094.75397
Hernando	FLA012070	Holiday Springs RV Resort WWTF	RIB	Discharge	194.6619633
Hernando	FLA012070	Holiday Springs RV Resort WWTF	Sprayfield	Discharge	0
Hernando	FLA012719	Shady Hills Elementary School WWTF	RIB	High	13.35052013
Pasco	FLA012741	Pasco County - Shady Hills Subregional WWTF	Reuse (golf, resi)	Medium	293013.9162
Pasco	FLA012748	Country Village MHP WWTF	RIB	Medium	87.53549638
Pasco	FLA012756	Big Oaks RV Park WWTF	RIB	High	66.80682413
Pasco	FLA012819	East Lake Landings MHP WWTF	RIB	High	268.8182339
Pasco	FLA012831	Travelers Rest WWTF	RIB	High	1145.906526
Pasco	FLA012831	Travelers Rest WWTF	Sprayfield	High	203.2149505
Pasco	FLA012832	Aripeka West MHP	RIB	Discharge	104.4095179
Hernando	FLA017033	Lakewood Retreat WWTF	Sprayfield	Medium	122.6042961
Hernando	FLA017223	Hernando Airport Subregional WWTF	RIB	High	11069.81546
Hernando	FLA186830	Eckerd Youth Challenge Program WWTF	Sprayfield	High	78.79736782



WWTFs

- For the entire Springs Coast region, 84 domestic wastewater facilities and discharge sites were evaluated to determine contributions to groundwater loading.
- Rapid infiltration basin and sprayfield applications are the most common disposal method across BMAP areas.
- The results of the NSILT analysis for all WWTFs and disposal types in all four springsheds are presented in the table below.

Springshed	Facilities	Est. Load to Surface (lbs-N/yr)	Est. Load to Groundwater (lbs-N/yr)
Kings Bay	23	71,397	36,645
Homosassa	22	8,544	3,382
Chassahowitzka	8	33,617	17,972
Weeki Wachee	31	415,919	79,729




OSTDS

- Florida Water Management Inventory data was used to estimate the number of OSTDS (updated 2021-2022).
- Parcels identified as "known septic," "likely septic," and "somewhat likely septic" were considered to have one septic system per parcel.
- Estimated load per tank is based on multiplying the average persons per household (2020 U.S. Census data) by loading per person, which is estimated to be 10 lbs/yr (Armstrong 2015).
- Credited enhanced nutrient reducing OSTDS with a 50% reduction in TN loading from the existing condition.
- Loading from OSTDS is estimated to attenuate at 30%.




OSTDS


Kings Bay/ Crystal River



Homosassa/ Chassaowitzka Springs



Weeki Wachee



*Density is per 300-meter by 300-meter grid cell.



OSTDS

The results of the NSILT analysis for OSTDS in all four springsheds are presented in the table below.

Springshed	Number of OSTDS	Number of enhanced OSTDS	Est. Load to Drain Fields (lbs-N/yr)	Est. Load to Groundwater (lbs-N/yr)
Kings Bay	30,256	638	673,211	413,555
Homosassa	16,947	56	364,522	215,178
Chassahowitzka	5,598	199	132,376	81,452
Weeki Wachee	44,726	1,114	1,048,663	641,621



UTF

- UTF loading was evaluated separately for single family residential, non-single family residential and sports turfgrass.
- Sports turfgrass loading includes the application of fertilizers to both sports fields and golf courses.
- Loading from UTFs are estimated to attenuate at 70%.



Source: Pexels



UTF

SINGLE FAMILY RESIDENTIAL

Single Family Residential Fertilization:

- 2021 Florida Department of Revenue Cadastral information was used to determine size and value of single-family residential parcels.
- Assumed 27.8% of parcel as impervious area (Tilley & Slonecker, 2006).
- Assumed 1 acre maximum for fertilized area for single family residential parcels greater than 1 acres in these springsheds.
- Assumed likelihood to fertilize rate is based on home value. Three tiers of home value considered.
- Fertilizer application rates are informed by local survey information and the Green Industries Best Management Practice (GIBMP) manual.

Lawn Care Source	
Service	32%
Self	68%

Self Application Rates	Percent	Application Rate (lbs/N/ac/applications)	Number of Applications
BMP	15%	26.136	5.00
Label	74%	34.78	2.98
None	11%	0.00	0.00



UTF

SINGLE FAMILY RESIDENTIAL

The results of the NSILT analysis for Single Family Residential UTF in all four springsheds are presented in the table below.

Springshed	Est. Load to Surface (lbs-N/yr)	Est. Load to Groundwater (lbs-N/yr)
Kings Bay	642,269	168,375
Homosassa	309,218	78,814
Chassahowitzka	177,242	46,716
Weeki Wachee	866,028	222,374



UTF

NON-SINGLE FAMILY RESIDENTIAL

- WMD land use/land cover data was used to estimate non-single family residential UTF application acreage estimates.
- Land use data year is dependent on the WMD with all data updated between 2019 and 2022.
- Fifteen land cover codes were assumed to be likely to receive fertilizer.

WMD Land Cover Code	Percent Impervious	Percent of Pervious area Receiving Fertilizer
1220: Medium Density, Mobile Home Units	32.6%	17.7%
1230: Medium Density, Mixed Units (Fixed and Mobile Home Units)	32.6%	15.4%
1320: High Density, Mobile Home Units	44.4%	20.7%
1330: Multiple Dwelling Units, Low Rise	44.4%	27.8%
1340: High Density, Multiple Dwelling Units, High Rise (Four Stories or More)	44.4%	32.8%
1400: Commercial and Services	72.2%	31.3%
1411: Shopping Centers	72.2%	31.3%
1480: Cemeteries	8.3%	42.2%
1700: Institutional	34.4%	43.3%
1710: Educational	30.3%	60.6%
1720: Religious	39.9%	37.7%
1740: Medical and Health Care	72.2%	33.8%
1750: Governmental	35.4%	41.0%
1850: Parks and Zoos	12.5%	44.9%
1860: Community Recreational Facilities	12.5%	59.8%

- Impervious area was estimated using a United States Geological Survey study (Tilley & Slonecker, 2006). Percent impervious area was dependent on the land use category.
- The area of pervious surface expected to receive fertilizer was evaluated by local land cover data. The percentage of pervious area evaluated to be bare ground or grass surface was applied to the estimated pervious areas of the 15 land cover codes to then estimate the area expected to receive fertilization.



UTF

NON-SINGLE FAMILY RESIDENTIAL

- Estimated fertilizer application rates for non-single family residential fertilized areas were based on an evaluation of the GIBMP Manual.
- It is estimated that fertilizer is applied at a rate of 3 lb-N/1,000 ft² to fertilized turfgrass in the region.
- The results of the NSILT analysis for non-single family residential UTF in all four springsheds are presented in the table below.

Springshed	Est. Load to Surface (lbs-N/yr)	Est. Load to Groundwater (lbs-N/yr)
Kings Bay	53,212	13,042
Homosassa	33,444	8,143
Chassahowitzka	20,110	5,237
Weeki Wachee	68,165	17,686



STF

STF is a combination of golf course and other sports turfgrass areas:

- Previous NSILT estimates of other STF areas were used in this evaluation to estimate loading from this source.
- Fertilizer application rates and area from the previous NSILT were used to estimate current nutrient loading where information was available.

Thirty-one golf courses operating within the springsheds were reviewed.

- Where site specific data was not available, golf course application rates were updated based on a study of regional golf course practices published by HortTechnology (Shaddox, et al., 2023).

Golf Course Study Rate		
Application Rate	95.832	lb-N/ac
% fertilized	100%	



Source: Pexels



STF

BMAP	County	Golf Course	Regional Rate or Site Specific Rate?	Land Area (Ac)				N Input (lb-N/yr)			
				High	Medium	Low	Discharge	High	Medium	Low	Discharge
KING	Citrus	Citrus Hills	Regional	284				27,216	-	-	-
KING	Citrus	Citrus Springs	Regional	241				23,096	-	-	-
KING	Citrus	Skyview	Regional	203				19,454	-	-	-
KING	Citrus	Twisted Oaks	Regional	134				12,841	-	-	-
KING	Citrus	Plantation Inn	Regional		198			-	18,975	-	-
KING	Citrus	Black Diamond	Site Specific	544				1,130			-
KING	Citrus	Lakeside	Site Specific	185				2,483			-
CHHO	Citrus	Sugarmill Woods Country Club	Regional	142				13,576	-	-	-
CHHO	Citrus	Inverness Golf and Country Club	Regional	123	5			11,776	492	-	-
CHHO	Citrus	Point O'Woods Golf Club	Regional	20				1,879	-	-	-
CHHO	Citrus	Cypress/Oak/Pine Courses - Sugarmill Woods Country Club	Regional	119.2	0.1			11,420	8	-	-
CHHO	Citrus	Southern Woods Golf Club (Citrus National Golf Club)	Regional	153				14,662	-	-	-
CHHO	Citrus	Sweet Swing Driving Range	Regional		8			-	767	-	-
CHHO	Hernando	Brooksville Country Club	Regional	182				17,441	-	-	-
CHHO	Hernando	Cabot Citrus Farms Golf Club	Regional	577	154			55,291	14,751	-	-
WEEK	Hernando	Hernando Oaks Golf and Country Club	Regional	139				13,321	-	-	-
WEEK	Hernando	Oak Hills Golf Club	Regional	184				17,676	-	-	-
WEEK	Hernando	Rivard Golf Club	Regional	95				9,147	-	-	-
WEEK	Hernando	Silverthorn Country Club	Regional	176				16,866			
WEEK	Hernando	Southern Hills Plantation Club	Regional	162				15,499	-	-	-
WEEK	Hernando	Spring Hill Golf and Country Club	Regional	159				15,194	-	-	-
WEEK	Hernando	Brookridge Country Club	Site Specific	85				32,844			-
WEEK	Hernando	Glen Lakes Golf and Country Club	Site Specific	86				602			-
WEEK	Hernando	Heather Golf and Country Club	Site Specific	55				4,875			-
WEEK	Hernando	High Point Golf Club	Site Specific	80				5,489			-
WEEK	Hernando	Timber Pines Country Club	Site Specific	215			43	684			4,121
WEEK	Pasco	Heritage Pines Community	Regional	194				18,591	-	-	-
WEEK	Pasco	Travelers Rest RV Resort and Golf Course	Site Specific					268			-



STF

The results of the NSILT analysis for STF, including golf courses, in all four springsheds are presented in the table below.

Springshed	Acres STF	Number of Golf Courses	Est. Load to Surface (lbs-N/yr)	Est. Load to Groundwater (lbs-N/yr)
Kings Bay	183	9	113,185	28,283
Homosassa	39	4	51,234	13,649
Chassahowitzka	99	4	100,152	25,178
Weeki Wachee	88	14	154,907	41,825



FF

- Florida Statewide Agricultural Irrigation Demand 9 (FSAID 9) data layer published in 2021 was analyzed to estimate acreage of all crop types within each recharge category.
- Application rates previously used in the NSILT were reviewed by the Florida Department of Agriculture and Consumer Services, WMDs, and University of Florida Institute of Food and Agricultural Science
- For all crops besides pasture and nurseries, loading to land surface was calculated by multiplying the acreage of a given crop type by the estimated fertilizer application rate.
- Loading from FF is estimated to attenuate at 80%.



FF


Nursery Crops	Fertilizer Application Rate (lb-N/acre)	Effective Application Rate (lb-N/acre)
Asparagus Fern	90	21.6
Aspidistra	90	21.6
Container Nursery	150	36
Coontie Fern	90	21.6
Fern	90	21.6
Field Nursery	90	21.6
Leatherleaf	90	21.6
Liriope	90	21.6
Nurseries and Vineyards	90	21.6
Nursery	90	21.6
Ornamentals	90	21.6
Pittosporum	90	21.6
Timber Nursery	50	12
Tree Nurseries	90	21.6

Crop	Fertilizer Application Rate (lb-N/acre)	Effective Application Rate (lb-N/acre)
Grass Pasture	80	16
Horse Farms	50	10
Improved Pastures	50	10
Pasture	50	10

- For nurseries that use containers:
 - A reduction of 20% of the FSAID 9 land area was made to account for plant spacing.
 - A reduction of 70% of loading was estimated to account fertilizer being applied to containers that hold the nutrients longer.
- Adjustment to fertilizer application rates were made to pastureland that utilize field rotation.
 - Fertilizer was estimated to be applied to 20% of pasturelands annually.




FF KINGS BAY



Kings Bay			
Crop	Total Acres	Loading to Land Surface (lb-N/year)	Loading to Groundwater (lb-N/year)
Grass Pasture	7,452	119,226	19,923
Cropland Pastureland	1,863	93,135	15,710
Hay	242	43,528	7,238
Specialty Farms	349	10,484	1,887
Blueberries	38	3,775	679
Improved Pastures	112	1,122	202
Row Crops	17	1,002	100
Field Nursery	23	502	90
Nursery	14	294	53
Pasture	21	206	37



FF HOMOSASSA-CHASSAHOWITZKA




Homosassa			
Crop	Total Acres	Loading to Land Surface (lb-N/year)	Loading to Groundwater (lb-N/year)
Grass Pasture	16,334	261,346	45,932
Cropland Pastureland	2,957	147,852	23,737
Melons	453	67,964	12,234
Hay	288	51,917	9,344
Other Hay Non Alfalfa	184	33,047	5,585
Dry Beans Tomatoes Spring	62	12,362	2,225
Specialty Farms	374	11,226	1,977
Blueberries	137	13,736	1,866
Strawberries	92	13,869	1,812
Citrus	79	11,037	1,266

Chassahowitzka			
Crop	Total Acres	Loading to Land Surface (lb-N/year)	Loading to Groundwater (lb-N/year)
Grass Pasture	6,782	108,505	18,616
Cropland Pastureland	1,709	85,457	14,450
Other Hay (Non Alfalfa)	312	56,143	10,047
Hay	138	24,801	4,464
Pasture	1,684	16,840	2,985
Specialty Farms	411	12,316	2,122
Field Nursery	438	9,458	1,702
Citrus	36	5,085	915
Improved Pastures	259	2,589	466
Blueberries	22	2,214	399



FF WEEKI WACHEE



Weeki Wachee			
Crop	Total Acres	Loading to Land Surface (lb-N/year)	Loading to Groundwater (lb-N/year)
Grass Pasture	25,662	410,588	60,389
Cropland Pastureland	3,938	196,925	30,017
Other Hay (Non Alfalfa)	606	109,070	19,381
Hay	622	111,953	13,050
Blueberries	442	44,220	7,141
Specialty Farms	672	20,172	3,176
Improved Pastures	1,371	13,713	1,748
Citrus	92	12,901	1,416
Nursery	434	9,384	967
Pasture	315	3,148	567



FF

The results of the NSILT analysis for FF in all four springsheds are presented in the table below.

Springshed	Est. Load to Surface (lbs-N/yr)	Est. Load to Groundwater (lbs-N/yr)
Kings Bay	273,328	45,930
Homosassa	631,905	108,876
Chassahowitzka	320,023	56,274
Weeki Wachee	946,876	139,819



LW

- Livestock populations were estimated using 2017 U.S. Department of Agriculture (USDA) census of agriculture data. USDA population estimates are provided by county.
- FSAID 9 land use was used to evaluate the proportion of livestock land within a BMAP and adjust USDA population estimates.
- Waste factors were multiplied by the estimated animal population to calculate LW loading.
- Loading from LW is estimated to attenuate at 90%.

Livestock Type	Waste Factor (lb-N/day)
Beef Cattle	0.337
"Other" Cattle	0.31
Calves	0.068
Dairy Cows	0.977
Donkeys	0.1
Horses	0.273
Chicken, Broilers	0.002
Chicken, Layers	0.003
Goats	0.035
Hogs	0.19
Sheep	0.198
Turkeys	0.006



LW

Kings Bay			
Livestock Type	Total Head Count	Loading to Land Surface (lb-N/year)	Loading to Groundwater (lb-N/year)
Beef Cattle	2,005	246,602	21,797
"Other" Cattle	22	2,520	223
Calves	1,092	13,583	1,201
Donkeys	81	2,965	265
Horses	590	58,773	5,259
Chicken, Broilers	18	13	1
Chicken, Layers	2,156	2,361	211
Goats	749	9,562	856
Hogs	152	10,547	944
Sheep	294	21,195	1,897
Turkeys	72	158	14

Homosassa			
Livestock Type	Total Head Count	Loading to Land Surface (lb-N/year)	Loading to Groundwater (lb-N/year)
Beef Cattle	3,668	451,099	35,977
"Other" Cattle	62	7,066	589
Calves	2,008	24,991	1,995
Donkeys	92	3,356	290
Horses	619	61,694	5,313
Chicken, Broilers	15	11	1
Chicken, Layers	2,086	2,285	196
Goats	729	9,314	801
Hogs	145	10,105	868
Sheep	272	19,695	1,690
Turkeys	89	193	17

Weeki Wachee			
Livestock Type	Total Head Count	Loading to Land Surface (lb-N/year)	Loading to Groundwater (lb-N/year)
Beef Cattle	12,370	1,521,536	115,942
"Other" Cattle	174	19,740	1,546
Calves	6,755	84,053	6,407
Donkeys	188	6,872	526
Horses	1,512	150,701	11,376
Chicken, Broilers	282	207	15
Chicken, Layers	684	749	63
Goats	799	10,217	783
Hogs	126	8,785	677
Sheep	332	23,974	1,810
Turkeys	179	394	30

Chassahowitzka			
Livestock Type	Total Head Count	Loading to Land Surface (lb-N/year)	Loading to Groundwater (lb-N/year)
Beef Cattle	4,901	602,982	53,001
"Other" Cattle	162	18,252	1,604
Calves	2,726	33,928	2,982
Donkeys	134	4,882	429
Horses	705	70,277	6,180
Chicken, Broilers	0	0	0
Chicken, Layers	1,606	1,759	155
Goats	584	7,461	656
Hogs	106	7,295	642
Sheep	156	11,295	993
Turkeys	157	344	30



DAIRIES

- Non-Confined Animal Feeding Operation dairy information was provided by Florida Department of Agriculture and Consumer Services, including information on herd size, waste handling practices and animal confinement.
- If a dairy herd was identified as grazed, it was estimated that they would be confined for 15% of the time to account for time in the milking parlors.
- A waste factor of 0.36 lb-N/day for dairy cows and 0.15 lb-N/day for non-milking cows was assumed.
- Annual loading was estimated by multiplying the number of cows by the daily waste factor multiplied by 365 days per year multiplied by application loss coefficients based on waste handling practices.
- Dairy waste was estimated to attenuate at 50% for the BMAP.



LW

The results of the NSILT analysis for livestock waste in all four springsheds are presented in the table below.

Springshed	Est. Load to Surface (lbs-N/yr)	Est. Load to Groundwater (lbs-N/yr)
Kings Bay	368,278	32,668
Homosassa	642,973	81,944
Chassahowitzka	755,071	66,674
Weeki Wachee	1,827,226	139,175



BIOSOLIDS

- Biosolids application quantity estimates were derived from calculating the average application quantity reported from 2018 to 2022, where data was available.
- Application quantities are provided in tons of material, it was assumed biosolid material has an estimated nitrogen content of approximately 5%.
- Loading to land surface was calculated by multiplying the average application quantity for the period of record by the estimated nitrogen content of 5%.
- Loading from biosolids is estimated to attenuate at 50%.


Springshed	Est. Load to Surface (lbs-N/yr)	Est. Load to Groundwater (lbs-N/yr)
Kings Bay	12,920	5,782
Chassahowitzka	20,096	9,043
Weeki Wachee	28,618	12,878



DRAFT NSILT LOADING RESULTS

KINGS BAY/CRYSTAL RIVER

Kings Bay	
Source	Annual Loading (lb-N/year)
Atmospheric Deposition	69,099
Wastewater Treatment Facilities	36,645
OSTDS	413,555
Urban Turfgrass Fertilizer	181,417
Sports Turfgrass Fertilizer	28,283
Farm Fertilizer	45,930
Livestock Waste	32,668
Biosolids	5,782
Total	813,379






DRAFT NSILT LOADING RESULTS

HOMOSASSA/CHASSAHOWITZKA

Homosassa	
Source	Annual Loading (lb-N/year)
Atmospheric Deposition	70,808
Wastewater Treatment Facilities	3,382
OSTDS	215,178
Urban Turfgrass Fertilizer	86,957
Sports Turfgrass Fertilizer	13,649
Farm Fertilizer	108,876
Livestock Waste	81,944
Total	580,794

Homosassa Spring Group (2023)






DRAFT NSILT LOADING RESULTS

HOMOSASSA/CHASSAHOWITZKA

Chassahowitzka	
Source	Annual Loading (lb-N/year)
Atmospheric Deposition	43,944
Wastewater Treatment Facilities	17,972
OSTDS	81,452
Urban Turfgrass Fertilizer	51,953
Sports Turfgrass Fertilizer	25,178
Farm Fertilizer	56,274
Livestock Waste	66,674
Biosolids	9,043
Total	352,490

Chassahowitzka Spring Group (2023)






DRAFT NSILT LOADING RESULTS

WEEKI WACHEE

Weeki Wachee Spring Group

Source	Annual Loading (lb-N/year)
Atmospheric Deposition	93,069
Wastewater Treatment Facilities	79,729
OSTDS	641,621
Urban Turfgrass Fertilizer	240,059
Sports Turfgrass Fertilizer	41,825
Farm Fertilizer	139,819
Livestock Waste	139,175
Biosolids	12,878
Total	1,388,175

Weeki Wachee Spring Group (2023)






BMAP MEETING

PUBLIC QUESTIONS PERIOD


Verbal Questions:

- We ask that questions and comments be limited to **3 minutes** so that we may hear from everyone.
- Please type in the chat if you would like to unmute yourself and ask a question or comment.



Written Comments:

- Submit written comments concerning today's meeting to BMAPProgram@FloridaDEP.gov.





BMAP UPDATES

SPRING VENT LOAD ANALYSIS

Calculated the current loading using the most recent 10 years of nitrate and discharge data.

Calculated the percent reduction using the TMDL and current loading.


Applied the spring vent percent reduction to the updated NSILT loading.

Estimate the total reduction needed to meet the TMDL.



BMAP UPDATES

SPRING VENT LOAD ANALYSIS



Source: Suwannee River WMD [What is a Spring?](#)



BMAP UPDATES

SPRING VENT LOAD ANALYSIS

BMAP	Draft Nitrate Loads (lb-N/yr)			
	Total Load at Spring Vent	TMDL Load	Required Reduction	Percent Required Reduction
Crystal River / Kings Bay ¹	453,400	259,009	194,392	43%
Homosassa Spring Group ¹	271,301	94,924	176,376	65%
Chassahowitzka Spring Group ¹	207,128	82,543	124,585	60%
Weeki Wachee Spring Group ²	308,909	95,265	213,644	69%

Upper 95% confidence interval - nitrate data from 2012 to 2022.

¹ TMDL target is 0.23 mg/L

² TMDL target is 0.28 mg/L



BMAP UPDATES

ALLOCATION AND REDUCTION APPROACH

- The percent reduction calculated from the spring vent analysis is applied to the estimated NSILT load to determine the overall required reduction needed in the basin.
- Each source will be evaluated for a reduction strategy.
- Responsible entities will receive an allocation based on the combined necessary reductions estimated by source for their area based on the NSILT loading.



BMAP UPDATES

ALLOCATION AND REDUCTION APPROACH

OSTDS

- Reduction strategy based on BMAP OSTDS requirements in Appendix D, with additional reductions based on actual loading from OSTDS.

WWTFs

- Reduction strategy is based on BMAP effluent requirements in the BMAP document and requirements in Florida law established 2021-2024.

Agriculture

- Reduction strategy based on:
 - BMP enrollment using a 15% reduction applied to FF load to groundwater.
 - BMP enrollment using a 10% reduction applied to (LW and dairies.
 - Any remaining agricultural reductions will be allocated to agricultural cooperative elements, which could include regional projects, cost-share practices and innovative technologies.

AD

- Anticipate reductions to be addressed by reductions from other sources and regional projects.



BMAP UPDATES

ALLOCATION AND REDUCTION APPROACH

UTF

- Apply the spring vent percent reduction to the total UTF load to groundwater and allocate to the applicable local governments.

STF

- Apply the spring vent percent reduction to the STF load to groundwater and allocate to the applicable governments.

Golf Courses

- Reduction based on requirement of all golf courses to submit information on implementation of BMPs and a nutrient management plan to address the nutrient loading.



BMAP UPDATES


FUTURE GROWTH

- **Domestic Wastewater Projections:**
 - Use wastewater to estimate future growth projections.
 - Start with population growth for each county from Bureau of Economic and Business Research:
 - 2040 Medium Growth Projections.
 - Proportion growth for each entity based on land area.
 - Distinguish the future population expected to be served by sewer versus those with OSTDS based on the most recent Florida Water Management Inventory for each BMAP county.
 - Use per person calculations to estimate future loads from WWTF and OSTDS.
- **Agriculture Projections:**
 - Exploring different tools to estimate future changes in agricultural acreage in the BMAPs to estimate changes in agricultural loading.




BMAP UPDATES

5-, 10-, AND 15-YEAR MILESTONES/REDUCTION SCHEDULE





SPRINGS BMAP UPDATES TIMELINE





RESOURCES

BMAP WEBSITE AND STORYMAPS

Basin Management Action Plans

[Home](#) » [Divisions](#) » [Division of Environmental Assessment and Restoration](#) » [Water Quality Restoration Program](#) » Basin M

Water Quality Restoration Program Quick Links

[Basin Management Action Plans \(BMAPs\)](#)

[Statewide Annual Report](#)

[Water Quality Grant Opportunities 2023-24](#)

[BMAP Public Meetings](#)

[Impaired Waters, TMDLs and Basin Management Action Plans Interactive Map](#)

[Tools and Guidance for](#)

What is a Basin Management Action Plan

A basin management action plan (BMAP) is a framework for water quality management that reduces pollutant loading through current and future projects and strategies. BMAPs establish permit limits on wastewater facilities, urban and agricultural best management practices, and other measures to achieve pollutant reductions established by a total maximum daily load (TMDL). BMAPs are developed in consultation with stakeholders and rely on local input and commitment for development and implementation. BMAPs are required by Department of Environmental Protection Secretarial Order and are listed in the

Water Quality Protection Grant

DEP has launched an [online grant portal](#) to provide eligible entities to apply for grant programs. Eligible entities include local governments, academic institutions, and non-profit organizations. The [application portal](#) opened July 5, 2023. Closing dates for individual grant programs are posted on the portal. Applicants are encouraged to



Florida Springs Basin Management Action Plans (BMAPs)

Welcome to the Florida Springs Basin Management Action Plan (BMAP) StoryMap

The springs BMAPs are developed with specific provisions for the protection and restoration of the state's Outstanding Florida Springs. This story map focuses on the springs-related BMAPs; for more details about other BMAPs or more information about the BMAP program in general, visit <https://floridadep.gov/bmaps>.

* The story map will display differently depending on the screen size and resolution being used. Story map best viewed in Chrome or Firefox.

Overview



1 (COPY) Nitrogen Source Inventory and Loading Tool...



2 (COPY) Statutes & Bills



3 (COPY) Crystal River - Kings Bay BMAP Story Map



4 (COPY) DeLeon Spring BMAP Story Map



5 (COPY) Gemini Springs BMAP Story Map



6 (COPY) Homosassa and Chassahowitzka Springs Group...




7 (COPY) Jackson Blue and Merritts Mill Pond BMAP Story Map



8 (COPY) Santa Fe River BMAP Story Map



9 (COPY) Silver Springs, Upper Silver Springs, and Rainbow...





RESOURCES

FUNDING OPPORTUNITIES



Florida Department of Environmental Protection
Funding Opportunities

[FloridaDEP.gov/Funding](https://www.floridadep.gov/Funding)





SUBSCRIBER PAGE

HOW TO CONTACT US



BMAPProgram@FloridaDEP.gov



REFERENCES

- Armstrong, J.H., (2015). Florida Onsite Sewage Nitrogen Reduction Strategies Study Final Report.
- Eller, K. T., & Katz, B. G. (2017). Nitrogen Source Inventory and Loading Tool: An integrated approach toward restoration of water-quality impaired karst springs. *Journal of Environmental Management*.
- Helgeson, T., & McNeal, M., (2009). A Reconnaissance-Level Quantitative Comparison of Reclaimed Water, Surface Water, and Groundwater.
- Shaddox, T.W., Unruh, B.J., Johnson, M.E., Brown, C.D., & Stacey, G. (2023). Nutrient Use and Management Practices on United States Golf Courses. HortTechnology.
- Tilley, J.S., & Slonecker, E.T. (2006). Quantifying the Components of Impervious Surfaces: U.S. Geological Survey Open-File Report 2006-1008.
- [GI-BMP Manual](#), Florida Institute of Food and Agricultural Science Extension, (ufl.edu).




BMAP MEETING

PUBLIC QUESTIONS PERIOD

Verbal Questions:

- We ask that questions and comments be limited to **3 minutes** so that we may hear from everyone.
- Please type in the chat if you would like to unmute yourself and ask a question or comment.



Written Comments:

- Submit written comments concerning today's meeting to BMAPProgram@FloridaDEP.gov.





THANK YOU

Chandler Keenan
Environmental Consultant
850-245-8555

Chandler.B.Keenan@FloridaDEP.gov

Samuel Hankinson, P.G.
Environmental Consultant
(850) 245-8086

Samuel.Hankinson@FloridaDEP.gov

Lauren Campbell, Ph.D.
Environmental Administrator
850-245-8083

Lauren.Campbell@FloridaDEP.gov

