
EcoSummary

Wekiwa Springs

5/20/03



Background

Florida DEP's Division of Recreation and Parks selected Wekiwa Springs, as well as springs from several other state parks, for biological and water quality monitoring. Data from these efforts will be used for documenting conditions within each park and for making resource protection decisions. Wekiwa Springs State Park is located in Orange County, Florida (Figure 1). The Wekiva River begins in the park and then flows northeast into the St. Johns River. Wekiwa Springs is a Floridan Aquifer-dominated system and is known for its recreational and aesthetic qualities.

Water chemistry samples were taken for ammonia, nitrate-nitrite, total Kjeldahl nitrogen, total phosphorus, color, turbidity, *Enterococci*, *Escherichia coli*, fecal coliform, and total coliform analysis; see Figure 2. Habitat assessment, qualitative periphyton sampling, and benthic invertebrate stream condition index (SCI) sampling were conducted.

Results

The nitrate-nitrite concentration (1.4 mg/L) at the sampling site was higher than those found in 95% of Florida streams (Figure 2). Elevated nitrite-nitrate levels in Wekiwa River are related to nitrogen loading in the recharge basin. The total Kjeldahl nitrogen and Ammonia concentrations were below quantitation limit (Figure 2). The total phosphorus concentrations (0.13 mg/L) were lower than those found in 60% of Florida streams.

The overall habitat assessment score (123) was in the optimal range (>120). *Enterococci* concentration (250 CFU/100 mL) was above the EPA single-sampling guideline value of 61 CFU/100 mL for "designated beach areas." Some species of *Enterococci* are known to be free-living in warm waters; the applicability of the above guideline in Florida waters is currently under investigation. *Escherichia coli* concentration (28 CFU/100 mL) was below the EPA designated beach guideline of 235 CFU/100 mL. Fecal and total coliform levels complied with Class III water quality standards. The stream condition index (SCI) score of 25 for Wekiwa Springs remained in the "good" range (scored 25 on 12/6/01, 4/17/02 and 10/22/02). The periphyton community in Wekiwa Springs was dominated by diatoms with some blue-green taxa. The particular taxa found were indicative of eutrophic conditions, which was not surprising given the continued nutrient enrichment in Wekiwa Springs.

In conclusion, elevated nitrate-nitrate levels in Wekiwa Springs continue to be the main stressor associated with a relatively low quality algal community.

Figure 1: Overview Map of the Wekiwa Springs Area




Figure 2: Data Table

Wekiwa Springs			
Sampling Date		5/20/2003	
Macroinvertibrate Parameters		Periphyton Parameters	
SCI	25	% Bacillariophyceae	87.5
SCI Evaluation	Good	% Cyanophyceae	11.8
Number of Taxa	31	% Chlorophyceae	0.6
EPT Index	1	Dominant taxon	Fragilariaceae
FL Index	12	Percent Dominant taxon	25.7
% Dominant Taxon	24	Physical-Chemical Data	
% Diptera	45.2	Habitat Assessment	123
Total number of Chironomidae	15	Specific Conductivity (umhos/cm)	345
% Filter-Feeders	9.6	Dissolved Oxygen (mg/L)	1.6
Chemistry Data		pH (SU)	7.2
Ammonia (mg/L)	0.018 I	Temperature (deg C)	23.6
Nitrate-Nitrite (mg/L)	1.4	Bacteria Parameters	
TKN (mg/L)	0.19 I	*Enterococci* (col/100mL)	250 Q
Total Phosphorus (mg/L)	0.13	*Escherichia coli* (col/100mL)	28 BQ
Color (PCU)	15	Fecal Coliforms (col/100 mL)	40 Q
Turbidity (NTU)	0.2	Total Coliforms (col/100mL)	520 Q

"I"- Below Quantitation Limit; "Q"- Information Only; "J"- Estimated Value

Figure 3. SCI Evaluations



FOR MORE INFORMATION, CONTACT:
Thomas Deck, FDEP Environmental Assessment Section
2600 Blair Stone Road MS#6511 Tallahassee, FL 32399-2400 (850) 245-8074 thomas.deck@dep.state.fl.us