


BIOASSAYS OF CHAMPION PAPER COMPANY

KRAFT DIVISION

CANTONMENT, ESCAMBIA COUNTY, FLORIDA

NPDES #FL0002526

SAMPLED 4/28/86

JULY 23, 1986

Biology Section
Division of Environmental Programs

BIOASSAYS OF CHAMPION PAPER COMPANY

KRAFT DIVISION

CANTONMENT, ESCAMBIA COUNTY, FLORIDA

NPDES #FL0002526

Biology Section
Bureau of Laboratories and
Special Programs
July 23, 1986

EXECUTIVE SUMMARY

Champion Paper Company, Cantonment, NPDES #FL0002526, on 29 April to 1 May, 1986

The sample of paper mill effluent collected from the company's Outfall #001 was not acutely toxic to the test organisms, Daphnia pulex and Notropis leedsi. No mortality occurred during the 48 hour static bioassays. The ammonia measured in the effluent (2.4 mg/l total and 0.064 mg/l unionized) was insufficient to have caused any toxicity. Since the sample was not acutely toxic, result of chemical analyses will not be presented in this report.

INTRODUCTION

The Champion Paper Company, formerly the St. Regis Paper Company, is located at the intersection of U.S. Highway 29 and State Road 184 in Cantonment, Escambia County (Fig. 1). According to the NPDES Permit (#FL0002526, issued in June 1979), this facility operates continuously, processing 1000 tons per day of pulpwood into paper products. Raw material required for this operation include: pulpwood (pine or hardwood), salt cake, lime, caustic soda, chlorine, salt, sodium chlorate, rosin size, alum, sulfuric acid, and cornstarch.


Figure 2 depicts the water flow through the Champion Plant in Cantonment. The following compounds are associated with the paper manufacturing process and may be found in the wastewater stream: pimaric acid, isoprimaric acid, sandaracopimaric acid, abietic acid, dehydroabietic acid, 3, 4, 5, trichloroquaiacol, 3, 4, 5, 6, tetrachloroquaiacol, mono-chloro-dehydroabietic acid (2 isomers), 9, 10 epoxystearic acid, hydrogen sulfide, methyl mercaptan, glycol inks, and sodium sulfide. Prior to discharge to the headwaters of Eleven Mile Creek, the wastewater is treated by a clarifier, a series of mechanically aerated lagoons and final settling in an additional lagoon. This system is designed for a 28 MGD wastewater flow. Champions Outfall #001 is located at State Road 186, approximately 2 miles from the plant site. Eleven Mile Creek eventually empties into the northern portion of Perdido Bay approximately 12 miles downstream of the plant outfall.

On 17 to 19 February, 1981, the Biology Section performed three separate static bioassays on an effluent sample from the Champion Paper Company, Outfall #001. The results of those tests indicated that the sample ws not toxic to the test organisms, Daphnia pulex, Simocephalus serrulatus, and Pimphales promelas.

To again determine the existence and extent of toxicity of this effluent, the Biology Section performed additional bioassays on samples of effluent collected from Champion Paper Company's Outfall #001 and from Eleven Mile Creek downstream of the outfall boil. The tests were performed from 20 to 23 August, 1985. The results of those tests indicated that both samples were acutely toxic to Notropis leedsi. Ammonia was believed to have been mostly responsible for this toxicity (FDER 1985a). As a result of this toxicity, the Northwest District requested that further bioassays be performed. A sample of effluent was collected from Champion's Outfall #001 on 28 April, 1986, and two 48 hour static bioassays were performed. The results of these latest tests are presented in this report. Those involved in performing the tests and in preparation of this report were: R. B. Frydenborg, M. L. Roll, and P. G. Weiland.



Figure 1. Portions of Escambia County, Florida, including the locations of the Champion facilit near Cantonment and the sampling point at S.R. 186.



FLOW UNITS - 1000 GAL/DAY

Figure 2. SCHEMATIC OF WATER FLOW
ST. REGIS PAPER COMPANY
CANTONMENT
ESCAMBIA COUNTY
FLORIDA
JUNE 20, 1978

METHODS AND MATERIALS

Test methods used to conduct these static bioassays are extensively described by Peltier and Weber (1985) and the FDER Biology Section (1985b). The organisms selected for this study include a species recommended by Peltier and Weber (1985), the U.S. EPA (1978), and the FDER Biology Section (1985). Additionally, the Biology Section maintains cultures of several species of organisms indigenous to the State of Florida. One of these indigenous species was also selected for testing. Life histories of these test organisms and maintenance procedures used to culture them are detailed below.

History and Maintenance of Bioassay Organisms

Daphnia pulex (water flea) - A breeding culture of Daphnia pulex was obtained from the EPA laboratory in Athens, Georgia, in March, 1979. This culture has been maintained since its arrival in 2 1/2 gallon all-glass aquaria, filled with unchlorinated DER well water, under a 16:8 hour light-dark cycle. Organisms are fed a liquified blend of Purina Trout Chow® and Cerophyl® twice daily, once daily on weekends, and are kept at a constant temperature of 22° ± 2°C. The aquaria are cleaned weekly by siphoning off half the water along with the detritus and excess organisms, and then refilling with fresh aerated well water. Additional measures are taken and/or new cultures started in the event of abnormal circumstances such as: declining populations, ephippia formation, or excessive fungal or bacterial growth.

Notropis leedsi (bannerfin shiner) - Approximately 40 adult bannerfin shiners were obtained from an indigenous population in the upper Suwannee River during October, 1981. The breeding stock is maintained in 30 and 125 gallon all-glass aquaria equipped with a flow-through water exchange system. Water -dark cycle and are fed Tetra-Min® and frozen Artemia (brin shrimp) twice daily, and once daily on weekends. Breeding and egg hatching are encouraged at 25°C. The eggs are laid between stacked red ceramic tiles separated by a 3mm bead of silicone. Most of the eggs adhere to the tiles and are easily removed to heated, aerated hatching tanks. The narrow space between the tiles eliminates egg predation by adult fish. New hatching tanks are started regularly to segregate fry into even age groups. Newly hatched fry are fed a liquefied blend of Purina Trout Chow® and Cerophyl® twice daily. After one week, feeding is augmented with newly hatched Artemia nauplii and finely crushed Tetra Min®. The fish used in this test were approximately 25 days old at the time of the test.

To ensure the uniform sensitivity of these test organisms to toxic substances, the Biology Section conducts monthly standard reference toxicant bioassays using sodium lauryl sulfate as the reference toxicant.

Test Methods

Methods used to perform acute toxicity bioassays are described by Peltier and Weber (1985) and the FDER Biology Section (1985b). Some of the methods described below are more specific than those of Peltier and Weber. These will be noted in more detail.

Sampling Methods and Test Dates

Two 48 hour static acute toxicity bioassays were performed on a sample of effluent collected from Champion's Outfall #001. The tests were performed from 1000 hours on 29 April to 1000 hours on 1 May, 1986.

The sample was collected by Mr. E. W. Chilvers of DER's Northwest District at 1343 hours on 28 April, 1986, using standard wastewater sampling procedures. Separate samples were obtained for chemical and biological testing. Samples for metal analyses were preserved with HNO_3. Samples for organic chemical analyses and for the bioassays were unpreserved. The samples were iced and transported to the DER laboratory in Tallahassee for analyses within 24 hours of collection.

Test Temperature

Upon arrival of the iced (4°C) bioassay samples in the Tallahassee laboratory, the containers were placed in warm water and the temperature of the samples allowed to rise to 20° $\pm$ 2°C. Although this temperature may be different than the temperature of the sample during collection, it is the temperature range to which the test organisms are acclimatized and is the range recommended in the above referenced bioassay protocols. This avoids stressful effects caused by exposure to unusually cold or warm samples.

Freshwater Tests

Freshwater for dilution of the effluent sample was obtained from the same DER deep well that is used to culture the test organisms. The concentration series selected for this study (Table 1) is that recommended by APHA (1980).

Two replicates, of each concentration and a control, were used to test with the water flea, *Daphnia pulex*. Flint glass jars of 250 ml capacity were used as test chambers, and 200 ml of the appropriate test concentration was placed in each chamber. *Daphnia* were siphoned into a large culture dish to isolate the proper size and age organisms (neonates). To accomplish this, the water fleas were siphoned through a stainless steel screen. The trapped adults were then discarded. These neonates were then transferred by glass pipette, examined for injury, and were loaded 10 per chamber, resulting in 20 per test concentration. Water fleas were not fed during this test.

Two replicates, of each concentration and a control, were used for the Notropis leedsi (bannerfin shiner) bioassay. The fish were transferred individually or by twos, into wide mouth quart jars containing 500 ml each of the appropriate test concentration, and were loaded 5 per chamber, resulting in 10 per concentration. Fish were not fed during this test.

Test Parameters

Dissolved oxygen (YSI Model 57 oxygen meter), pH (Corning Model 7 pH meter with Orion Ross® combination electrode), temperature (NBS calibrated glass thermometer), and the number of live organisms were recorded for each test concentration at 0 hours, 24 hours, and 48 hours. Total alkalinity and total hardness (measured as $CaCO_3$ using HACH Chemical Company reagents) were recorded at the beginning of the test for the controls and the 100% effluent concentration. The specific conductance (YSI Model 33 S-C-T meter) of each test concentration was recorded at the beginning and end of the test. All instruments were calibrated daily according to manufacturer's recommendations. Ammonia concentrations, including the total and unionized fraction (expressed in mg/l as N and as NH_3, respectively), were recorded in the controls and the 100% effluent test concentration at the beginning of the test. Total ammonia concentrations were determined using an Orion® specific ion probe and an Orion Ionalyzer®. The unionized fraction was then calculated based on the total ammonia concentration, and the pH, temperature, and salinity of the sample. These values were recorded on the bioassay data forms and are presented in Tables 2 and 3.

Table 1. Concentration Series and Volumes Used for These Static Bioassays.

Daphnia pulex (water flea)

Concentration	Dilution Water	Effluent	Total Volume
Control	200.0 ml	--------	200 ml
10.0%	180.0 ml	20.0 ml	200 ml
18.0%	164.0 ml	36.0 ml	200 ml
32.0%	136.0 ml	64.0 ml	200 ml
56.0%	88.0 ml	112.0 ml	200 ml
100.0%	--------	200.0 ml	200 ml

Notropis leedsi (bannerfin shiner)

Concentration	Dilution Water	Effluent	Total Volume
Control	500 ml	--------	500 ml
10.0%	450 ml	50 ml	500 ml
18.0%	410 ml	90 ml	500 ml
32.0%	340 ml	160 ml	500 ml
56.0%	220 ml	280 ml	500 ml
100.0%	------	500 ml	500 ml

Table 2. Data recorded during 48 hour, Daphnia pulex, static acute toxicity bioassays of Champion Paper Company, Cantonment, Escambia County, Florida, NPDES #FL0002526, on 29 April to 1 May, 1986.

Concentration or %	Number of Live Organisms			Dissolved Oxygen (mg/l)			pH			Temperature (°C)			Conductivity (umhos)	
	0	24	48	0	24	48	0	24	48	0	24	48	0	48
Control	10	10	10	8.8	9.1	8.9	8.1	7.9	7.7	22.3	20.0	21.4		
Control	10	10	10	8.8	9.1	8.9	8.1	7.9	7.7				225	260
10.0%	10	10	10	8.8	9.1	9.0	8.0	7.9	7.8	22.2	19.7	21.3		
10.0%	10	10	10	8.8	9.1	9.0	8.0	7.9	7.8				338	365
18.0%	10	10	10	8.8	9.0	9.0	8.0	7.9	7.8	22.2	19.6	21.2		
18.0%	10	10	10	8.8	9.0	9.0	8.0	7.9	7.8				512	440
32.0%	10	10	10	8.7	9.0	9.0	7.9	7.9	7.8	22.1	19.5	21.2		
32.0%	10	10	10	8.7	9.0	9.0	7.9	7.9	7.8				620	700
56.0%	10	10	10	8.5	8.6	8.2	7.8	7.9	7.8	21.8	19.5	21.2		
56.0%	10	10	10	8.5	8.6	8.2	7.8	7.9	7.8				905	1000
100.0%	10	10	10	8.0	8.1	7.4	7.7	7.8	7.9	21.5	20.0	21.3		
100.0%	10	10	10	8.0	8.1	7.4	7.7	7.8	7.9				1330	1450

Control Water

NH_3(as NH_3)(mg/l): 0

Total ammonia (as N)(mg/l): 0

Total Alkalinity (mg/l $CaCO_3$): 108

Total Hardness (mg/l $CaCO_3$): 115

100% Effluent

NH_3(as NH_3)(mg/l): 0.064

Total ammonia (as N)(mg/l): 2.4

Total Alkalinity (mg/l $CaCO_3$): 195

Total Hardness (mg/l $CaCO_3$): 145

Table 3. Data recorded during 48 hour, Notropis leedsi, static acute toxicity bioassay of Champion Paper Company, Cantonment, Escambia County, Florida, NPDES #FL0002526, on 29 April to 1 May, 1986.

Concentration or %	Number of Live Organisms			Dissolved Oxygen (mg/l)			pH			Temperature (°C)			Conductivity (umhos)	
	0	24	48	0	24	48	0	24	48	0	24	48	0	48
Control	5	5	5	8.5	8.9	8.9	7.6	7.8	7.7	22.7	18.8	21.5		
Control	5	5	5	8.5	8.9	8.9	7.6	7.8	7.7				212	215
10.0%	5	5	5	8.6	8.9	9.0	7.6	7.8	7.8	22.5	18.8	21.3		
10.0%	5	5	5	8.6	8.9	9.0	7.6	7.8	7.8				320	318
18.0%	5	5	5	8.6	8.9	8.9	7.6	7.8	7.8	22.2	18.7	21.1		
18.0%	5	5	5	8.6	8.9	8.8	7.6	7.8	7.8				405	400
32.0%	5	5	5	8.5	8.7	8.7	7.7	7.8	7.8	21.9	18.7	21.0		
32.0%	5	5	5	8.5	8.7	8.7	7.7	7.8	7.8				620	700
56.0%	5	5	5	8.3	8.1	8.0	7.7	7.8	7.8	21.8	18.9	21.0		
56.0%	5	5	5	8.3	8.1	8.0	7.7	7.8	7.8				905	910
100.0%	5	5	5	7.9	7.3	6.7	7.7	7.8	7.8	21.1	19.2	21.4		
100.0%	5	5	5	7.9	7.3	6.7	7.7	7.8	7.8				1350	1400

Control Water

NH_3(as NH_3)(mg/l): 0

Total ammonia (as N)(mg/l): 0

Total Alkalinity (mg/l $CaCO_3$): 108

Total Hardness (mg/l $CaCO_3$): 115

100% Effluent

NH_3(as NH_3)(mg/l): 0.061

Total ammonia (as N)(mg/l): 2.4

Total Alkalinity (mg/l $CaCO_3$): 195

Total Hardness (mg/l $CaCO_3$): 145

RESULTS AND DISCUSSION

The sample of effluent collected from Champion Paper Company's Outfall #001 was not acutely toxic to the test organisms, Daphnia pulex and Notropis leedsi. No mortality occurred during the 48 hour static bioassays. The ammonia measured in the effluent (2.4 mg/l total and 0.064 mg/l unionized) was insufficient to have caused any toxicity. Since the sample was not acutely toxic, results of chemical analyses will not be presented in this report.

LITERATURE CITED

American Public Health Association. 1980. Standard Methods for the Examination of Water and Wastewater. 15th ed. 1134 pp.

Florida Department of Environmental Regulation, Biology Section. 1981. Bioassays of St. Regis Paper Companny, Kraft Company Penscola Mill, Escambia County, Florida. NPDES #FL0002526. FDER.

_____. 1985a. Bioassays of Champion Paper Company, Draft Division, Cantonment, Escambia County, Florida. NPDES #FL0002526. FDER.

_____. 1985b. Quality Assurance Manual for Performing Acute Toxicity Tests (Second Edition). FDER.

Peltier, W. and C. I. Weber. 1985. Methods for measuring the acute toxicity of effluents to aquatic organisms (Third Edition). Environmental Research Laboratory. Athens, Georgia. EPA-600/4-85-012. 217 pp.