


EcoSummary

Wekiwa Springs

5/9/2007



Background

Florida DEP's Division of Recreation and Parks selected Wekiwa Springs for biological and water quality monitoring. Data from these efforts will be used for documenting conditions within the park and for making resource protection decisions. Wekiwa Springs State Park is located in Orange County, Florida (Figure 1). The Wekiva River begins in the park and then flows northeast into the St. Johns River. Wekiwa Springs is a Floridan Aquifer-dominated system and is known for its recreational and aesthetic qualities.

Water chemistry samples were taken for ammonia, nitrate-nitrite, total Kjeldahl nitrogen (TKN), total phosphorus, color, turbidity, *Enterococci*, *Escherichia coli*, fecal coliform, and total coliform analysis; see Figure 2. Habitat assessment and benthic invertebrate Stream Condition Index (SCI) sampling were also conducted.

Figure 1: Map of the Wekiwa Springs Area



Results

The nitrate-nitrite concentration (0.71 mg/L) at the sampling site was higher than the values found in 90% of typical Florida streams (0.64 mg/L or less). Elevated nitrate-nitrite levels in Wekiwa Springs are related to nitrogen loading in the recharge basin. The TKN concentration was below the practical quantitation limit, and ammonia was undetected. The total phosphorus concentration (0.13 mg/L) was average for typical Florida streams.

Enterococci and *Escherichia coli* concentrations complied with the EPA single-sample guideline values of 61 CFU/100 mL and 235 CFU/100 mL, respectively, for "designated beach areas." Fecal and total coliform levels complied with Class III water quality standards.

The thresholds for interpreting the Stream Condition Index values have recently been revised according to the EPA Biological Condition Gradient (BCG) conceptual model. In addition, the laboratory methodology used to evaluate the macroinvertebrate samples taken for the SCI was also revised to reduce uncertainty associated with SCI results. In the SCI 2004 method, a single group of 100 individuals was sub-sampled, identified, and scored, while in the new SCI 2007 method, two separate groups of 150 individuals are analyzed, and the two resulting SCI scores are averaged. The SCI 2007 produces a more robust evaluation since the decision thresholds are based on the BCG, and the examination of two separate aliquots reduces error. Note that the sampling method and the ten metrics used for calculating the score remain exactly the same. The SCI 2007 went into effect August 1, 2006. In Figure 3, the new thresholds have been retroactively applied to the scores acquired before this date.


The SCI score (41) was in the "healthy" category, and the Habitat Assessment score (133) was in the "optimal" range. In conclusion, while elevated nitrate-nitrite levels appeared to be the main issue of concern in Wekiwa Springs, biological health was good..

Figure 2: Data Table

WEKIWA SPRINGS			
Sampling Date	5/9/2007	**Field Data**	
Macroinvertebrate Data		Habitat Assessment Score	133
SCI Score	41 A	Temperature (deg. C)	23.7
SCI Evaluation	Healthy	pH	7.5
Chemistry Data		Dissolved Oxygen (mg/L)	3.9
Turbidity (NTU)	0.25	Specific Conductance (umhos/cm)	342
Color (PCU)	5 U	**Bacteria Data**	
Ammonia (mg/L)	0.010 U	*Escherichia coli* (col/100 mL)	42 Q
Nitrate-Nitrite (mg/L)	0.71	*Enterococci* (col/100 mL)	48 Q
TKN (mg/L)	0.14 I	Fecal Coliforms (col/100 mL)	78 Q
Total Phosphorus (mg/L)	0.13	Total Coliforms (col/100 mL)	114 Q

"U"= Below Method Detection Limit; "I"= Below Practical Quantitation Limit; "Q"= Outside of holding time; "A"= Value reported is the mean of two or more determinations

Figure 3. SCI Scores Over Time



FOR MORE INFORMATION, CONTACT:
Emily Peffer, FDEP Environmental Assessment Section
2600 Blair Stone Road MS# 6511 Tallahassee, FL 32399-2400
(850) 245-8071 Emily.Peffer@dep.state.fl.us