


APPENDIX A:

Caloosahatchee River Estuary Tape Grass Restoration Project

(submitted June 2014)



Acreage
 Dense: 1292.40
 Moderate: 397.09
 Scattered: 322.95
 Total Acreage: 2012.44



Data Sources:
 2004 One Meter Ortho Imagery provided by
 USGS and NRCS for Lee County, FL
 Coverage polygons digitized from USGS
 2004 aerial

Map Created by Daniel Dickinson

Figure 2. Proposed SAV restoration areas of *Vallisneria americana* in the upper Caloosahatchee Estuary in Lee County, Florida using seeding, whole plants and exclosures.




Figure 3. Location of *Vallisneria americana* restoration plantings and exclosures upstream of S-79 “Franklin Locks” permitted in May 2012 and currently being monitored by FGCU researchers.

GPS Coordinates

Site 1: Latitude: 26°43'8.80"N; Longitude: 81°39'57.60"W

Site 2: Latitude: 26°43'7.80"N; Longitude: 81°39'15.60"W

Site 3: Latitude: 26°43'12.10"N; Longitude: 81°39'6.20"W

Site 4: Latitude: 26°43'8.70"N; Longitude: 81°38'54.10"W




Figure 4. Additional proposed *Vallisneria americana* planting sites in freshwater tributaries to the Caloosahatchee River in Lee County, FL east of I-75. West to east; Orange River, Owl Creek, Trout Creek, Telegraph Creek, and Bedman Creek



Figure 5. Herbivore exclusion cage (6m x 2m) being deployed May 2012 by FGCU researchers in the Caloosahatchee River and planted with *Vallisneria americana* (10 shoots/m²).



Figure 6. Successful tape grass, *V. americana* restoration area in Caloosahatchee River from pilot study conducted by FGCU comparing two types of herbivore exclosures (Ceilley 2009).



Hardened exclosure cage with anchoring by J Anderson and Aquatic Ecosystems Solutions.



Small hardened exclosure cages planted with tape grass and deployed in flowing stream habitats of Florida

References Used to Prepare this Proposal:

- Bortone, S. A. and R. K. Turpin. 2000. Tape grass life history metrics associated with environmental variables in a controlled estuary. In: Bortone, S.A. (ed.), *Seagrasses: Monitoring, Ecology, Physiology, and Management*. CRC Press, Boca Raton, Florida, pp.65-79.
- Ceilley, D.W. 2009. Tape grass (*Vallisneria americana*) restoration pilot study: A comparison of two types of enclosures. Final Report to Lee County and the West Coast Inland Navigation District. Ft. Myers, FL.
- Ceilley, D.W., E. M. Everham, III, K.M. Ross, J.A. Ferlita and R.C. Young. 2011. Planting Tape Grass, *Vallisneria americana* in the freshwater Caloosahatchee River: A Dry Season Opportunity. Final Report to the South Florida Water Management District, W. Palm Beach, FL. 32 pp.
- Ceilley, D.W., I. Bartoszek, G. G. Buckner, and M.J. Schuman. 2003. Tape grass (*Vallisneria americana*) restoration feasibility study. Final Report to the South Florida Water Management District, 3300 Gun Club Rd. West Palm Bch, FL. 16 pp.
- Chamberlain, R. H. and P. H. Doering. 1998a. Freshwater inflow to the Caloosahatchee Estuary and the resource-based method for evaluation, p. 81-90. In S.F. Treat (ed.), *Proceedings of the 1997 Charlotte Harbor Public Conference and Technical Symposium* South Florida Water Management District and Charlotte Harbor National Estuary Program, Technical Report No. 98-02. Washington, D.C.
- Chamberlain, R. H. 2005. C-43 Basin Storage Reservoir Project - Caloosahatchee Estuary Hydrologic Evaluation Performance Measures. C-43 BSR Study Team Adopted Draft, June 28, 2005.
- Chamberlain, R. H. and P. H. Doering. 1998b. Preliminary estimate of optimum freshwater inflow to the Caloosahatchee Estuary: A resource-based approach, p. 121-130. In S.F. Treat (ed.), *Proceedings of the 1997 Charlotte Harbor Public Conference and Technical Symposium*. South Florida Water Management District and Charlotte Harbor National Estuary Program, Technical Report No. 98-02. Washington, D.C.
- Doering, P. H., R. H. Chamberlain, K. M. Donohue, and A. D. Steinman. 1999. Effect of salinity on the growth of *Vallisneria americana* Michx. from the Caloosahatchee Estuary, Florida. *Florida Scientist*. Vol. 62, No.2. pp. 89-105.
- Doering, P. H., R. H. Chamberlain, and D. E. Haunert. 2002. Using submerged aquatic vegetation to establish minimum and maximum freshwater inflows to the Caloosahatchee Estuary, Florida. *Estuaries* 25 (1343-1354).
- Doering, P. H., R. H. Chamberlain, and J. M. McMunigal. 2001. Effects of simulated saltwater intrusions on the growth and survival of wild celery, *Vallisneria americana*, from the Caloosahatchee Estuary (South Florida). *Estuaries*. Vol. 24. No. 6A, pp. 894-903.
- Fugate, D.C., and R.J. Chant, 2005, Near bottom stresses in a small highly stratified estuary, *Journal of Geophysical Research* 110, C03022, doi:10.1029/2004JC002563.
- Fugate, D.C., and C.T. Friedrichs, 2007, Lateral dynamics and associated transport of sediment in the upper reaches of a partially mixed estuary, Chesapeake Bay, USA, *Continental Shelf Research* 27, 679-698.
- Hauxwell, J., C.W. Osenberg, and T.K. Frazer. 2004. Conflicting management goals: manatees and invasive competitors inhibit restoration of a native macrophyte. *Ecological Applications*, 14(2), 2004, pp. 571-586 by the Ecological Society of America, Washington, D.C.

- Hoffacker, V.A. 1994. 1993 Caloosahatchee River Submerged Grass Observations. W.Report from Dexter Bender and Associates, Inc. to South Florida Water Management District, Fort Myers, FL.
- Hunt, H.L., Fugate, D.C., and Chant, R.J., 2009. Modeling bedload transport of juvenile bivalves: predicted changes in distribution and scale of postlarval dispersal, *Estuaries and Coasts* 32(6), 1090-1102.
- Hunt, M. J. 2003. An Ecological Model to Predict *V. americana* Densities in the Upper Caloosahatchee Estuary. In Proceedings of Greater Everglades Restoration Science Conference, Palm Harbor, FL. April 13-18.
- Hunt, M. J. and P. H. Doering. 2005. Significance of Considering Multiple Environmental Variables When Using Habitat as an Indicator of Estuarine Condition. In Bortone, S. A.(ed.), *Estuarine Indicators*, CRC Press, Boca Raton, FL, pp.221-227.
- Hunt, M. J., P. H. Doering, R. H. Chamberlain, and K. M. Haunert. 2003. Light and Salinity Stress for SAV Determined by Modeling and Experimental Work in the Oligohaline Zone of an Estuary. In Proceedings of Estuarine Research Federation, Seattle, WA. September.
- Kraemer, G. P., R. H. Chamberlain, P. H. Doering, A. D. Steinman, and M. D. Hanisak. 1999. Physiological responses of transplants of the freshwater angiosperm *Vallisneria americana* along a salinity gradient in the Caloosahatchee Estuary (Southwestern Florida). *Estuaries*. Vol. 22, No. 1 pp. 138-148.
- Mazzotti, F.J., L.G. Pearlstine, R. H. Chamberlain, M. J. Hunt, T. Barnes, K. Chartier, and D. DeAngelis. 2008. Stressor Response Model for Tape Grass (*Vallisneria americana*). CIR 1524, Department of Wildlife Ecology and Conservation, Florida Cooperative Extension Service, Institute of Food and Agricultural Sciences, University of Florida. 21 pp.
- Moore, K.A., E.C. Shields, and J.C. Jarvis. 2010. The role of habitat and herbivory on the restoration of tidal freshwater submerged aquatic vegetation populations. *Restoration Ecology* Vol.18, No. 4, pp. 596-604.
- RECOVER. 2004. CERP Monitoring and Assessment Plan: Part 1, Monitoring and Supporting Research. Restoration Coordination and Verification Team (RECOVER), c/o United States Army Corps of Engineers, Jacksonville District, Jacksonville, FL, and South Florida Water Management District, West Palm Beach, FL.
- Robbins, B.D. 2004. Habitat use of *Vallisneria americana* beds in the Caloosahatchee River: Final Report to the South Florida Water Management District. Mote Marine Lab. Tech Rpt; No.993.
- SCCF Marine Lab. 2012. Submersed vascular macrophyte restoration and monitoring in the Caloosahatchee: RESTORE Act Proposal submitted to CHNEP office 9/28/12.
- Smart, R. M., and J. D. Dorman. 1993. Latitudinal differences in the growth strategy of a submerged aquatic plant: ecotype differences in *Vallisneria americana*? *Bull.Ecol. Soc.Am.* 74 (Suppl.):439.
- South Florida Water Management District. 2003. Technical Documentation to Support Development of Minimum Flows and Levels for the Caloosahatchee River and Estuary, Status Update Report. May 2003.
- South Florida Water Management District. 2006. South Florida Environmental Report (Draft). Chapter 12 - Caloosahatchee River, Estuary, and Southern Charlotte Harbor.
- Titus, J. E. and M. S. Adams. 1979. Coexistence and the comparative light relations of the submerged macrophytes *Myriophyllum spicatum* L. and *Vallisneria americana* Michx. *Oecologia*. 40:273-286.
- Wilhere, G. F. 2002. Adaptive management in habitat conservation plans. *Conserv. Biol.* 16(1): 20-29.
- Xie, Y., S. Ana, X. Yoo, K. Xiaob, and C. Zhang. 2005. Short-time response in root morphology of *Vallisneria spiralis* to sediment type and water-column nutrient. *Aquatic Botany* 81: 85-96.

Bortone, S. A. and R. K. Turpin. 2000. Tape grass life history metrics associated with environmental variables in a controlled estuary. In: Bortone, S.A. (ed.), *Seagrasses: Monitoring, Ecology, Physiology, and Management*. CRC Press, Boca Raton, Florida, pp. 65-79.

Ceilley, D.W., I. Bartoszek, G. G. Buckner, and M.J. Schuman. 2003. Tape grass (*Vallisneria americana*) restoration feasibility study. Final Report to the South Florida Water Management District, 3300 Gun Club Rd. West Palm Bch, FL. 16 pp.

Ceilley, D.W. 2009. Tape grass (*Vallisneria americana*) restoration pilot study: A comparison of two types of exclosures. Final Report to Lee County and the West Coast Inland Navigation District. Ft. Myers, FL.

Ceilley, D.W., E. M. Everham, III, K.M. Ross, J.A. Ferlita and R.C. Young. 2011. Planting Tape Grass, *Vallisneria americana* in the freshwater Caloosahatchee River: A Dry Season Opportunity. Final Report to the South Florida Water Management District, W. Palm Beach, FL. 32 pp.

Chamberlain, R. H. 2005. C-43 Basin Storage Reservoir Project - Caloosahatchee Estuary Hydrologic Evaluation Performance Measures. C-43 BSR Study Team Adopted Draft, June 28, 2005.

Chamberlain, R. H. and P. H. Doering. 1998a. Freshwater inflow to the Caloosahatchee Estuary and the resource-based method for evaluation, p. 81-90. In S.F. Treat (ed.), *Proceedings of the 1997 Charlotte Harbor Public Conference and Technical Symposium*. South Florida Water Management District and Charlotte Harbor National Estuary

Program, Technical Report No. 98-02. Washington, D.C.

Chamberlain, R. H. and P. H. Doering. 1998b. Preliminary estimate of optimum freshwater inflow to the Caloosahatchee Estuary: A resource-based approach, p. 121-130. *In* S.F. Treat (ed.), Proceedings of the 1997 Charlotte Harbor Public Conference and Technical Symposium. South Florida Water Management District and Charlotte Harbor National Estuary Program, Technical Report No. 98-02. Washington, D.C.

Chamberlain, R. H. and P. H. Doering. 2004. Recommended Flow Distribution (the Caloosahatchee Estuary and the C-43 basin Storage Reservoir Project). Technical Memorandum, South Florida Water Management District.

18

Chamberlain, R. H., P. H. Doering, and K. M. Haunert. 2003. Preliminary assessment of water quality and material loading in the Caloosahatchee Estuary, FL. Poster Presentation, 17th Biennial Conference of the Estuarine Research Federation.

Chamberlain, R. H., D. E. Haunert, P. H. Doering, K. M. Haunert, and J. M. Otero. 1995. Preliminary estimate of optimum freshwater inflow to the Caloosahatchee Estuary, Florida. Technical report, South Florida Water Management District, W. Palm Bch. FL.

Doering, P. H. and R. H. Chamberlain 1998. Water quality in the Caloosahatchee Estuary, San Carlos Bay and Pine Island Sound. Proceedings of the Charlotte Harbor Public Conference and Technical Symposium; 1997 March 15-16; Punta Gorda, FL. Pp 229-240. Charlotte Harbor National Estuary Program Technical Report No. 98-02. 274 pp.

Doering, P. H., R. H. Chamberlain, K. M. Donohue, and A. D. Steinman. 1999. Effect of salinity on the growth of *Vallisneria americana* Michx. from the Caloosahatchee Estuary, Florida. *Florida Scientist*. Vol. 62, No.2. pp. 89-105.

Doering, P. H., R. H. Chamberlain, and D. E. Haunert. 2002. Using submerged aquatic vegetation to establish minimum and maximum freshwater inflows to the Caloosahatchee Estuary, Florida. *Estuaries* 25 (1343-1354).

Doering, P. H., R. H. Chamberlain, and J. M. McMunigal. 2001. Effects of simulated saltwater intrusions on the growth and survival of wild celery, *Vallisneria americana*, from the Caloosahatchee Estuary (South Florida). *Estuaries*. Vol. 24. No. 6A, pp. 894-903.

Hoffacker, V.A. 1994. 1993 Caloosahatchee River Submerged Grass Observations. W. Report from Dexter Bender and Associates, Inc. to South Florida Water Management District, Fort Myers, FL.

Fugate, D.C., and R.J. Chant, 2005, Near bottom stresses in a small highly stratified estuary, *Journal of Geophysical Research* 110, C03022, doi:10.1029/2004JC002563.

Fugate, D.C., and C.T. Friedrichs, 2007, Lateral dynamics and associated transport of sediment in the upper reaches of a partially mixed estuary, Chesapeake Bay, USA, *Continental Shelf Research* 27, 679-698.

Hunt, H.L., Fugate, D.C., and Chant, R.J., 2009. Modeling bedload transport of juvenile bivalves: predicted changes in distribution and scale of postlarval dispersal, *Estuaries and Coasts* 32(6), 1090-1102.

Hunt, M. J. 2003. An Ecological Model to Predict *V. americana* Densities in the Upper Caloosahatchee Estuary. In Proceedings of Greater Everglades Restoration Science Conference, Palm Harbor, FL. April 13-18.

Hunt, M. J. and P. H. Doering. 2005. Significance of Considering Multiple Environmental Variables When Using Habitat as an Indicator of Estuarine Condition. In Bortone, S. A. (ed.), *Estuarine Indicators*, CRC Press, Boca Raton, FL, pp.221-227.

Hunt, M. J., P. H. Doering, R. H. Chamberlain, and K. M. Haunert. 2003. Light and Salinity Stress for SAV Determined by Modeling and Experimental Work in the Oligohaline Zone of an Estuary. In Proceedings of Estuarine Research Federation, Seattle, WA. September.

Hunt, M. J., P. H. Doering, R. H. Chamberlain, and K. M. Haunert. 2004. Grass Bed Growth and Estuarine Condition: Is size a Factor worth Considering? In Conference Proceedings of the Southeastern Estuarine research Society, Harbor Branch Oceanographic Institution, Fort Pierce, FL. April 15-17.

Kraemer, G. P., R. H. Chamberlain, P. H. Doering, A. D. Steinman, and M. D. Hanisak. 1999. Physiological responses of transplants of the freshwater angiosperm *Vallisneria americana* along a salinity gradient in the Caloosahatchee Estuary (Southwestern

Florida). *Estuaries*. Vol. 22, No. 1 pp. 138-148.

Mazzotti, F.J., L.G. Pearlstine, R. H. Chamberlain, M. J. Hunt, T. Barnes, K. Chartier, and D. DeAngelis. 2008. Stressor Reponse Model for Tape Grass (*Vallisneria americana*). CIR 1524, Department of Wildlife Ecology and Conservation, Florida Cooperative Extension Service, Institute of Food and Agricultural Sciences, University of Florida. 21 pp.

Moore, K.A., E.C. Shields, and J.C. Jarvis. 2010. The role of habitat and herbivory on the restoration of tidal freshwater submerged aquatic vegetation populations. *Restoration Ecology* Vol.18, No. 4, pp. 596-604.

Ogden, J. C., S. M. Davis, and L. A. Brandt. 2003. Science Strategy for a Regional Ecosystem Monitoring and Assessment Program. In *Monitoring Ecosystems*, eds. D.E. Busch and J.C. Trexler, 135-163. Washington, DC: Island Press.

RECOVER. 2004. CERP Monitoring and Assessment Plan: Part 1, Monitoring and Supporting Research. Restoration Coordination and Verification Team (RECOVER), c/o United States Army Corps of Engineers, Jacksonville District, Jacksonville, FL, and South Florida Water Management District, West Palm Beach, FL.

Robbins, B.D. 2004. Habitat use of *Vallisneria americana* beds in the Caloosahatchee River: Final Report to the South Florida Water Management District. Mote Marine Lab. Tech Rpt; No. 993.

SCCF Marine Lab. 2012. Submersed vascular macrophyte restoration and monitoring in the Caloosahatchee: RESTORE Act Proposal submitted to CHNEP office 9/28/12.

Smart, R. M., and J. D. Dorman. 1993. Latitudinal differences in the growth strategy of a submerged aquatic plant: ecotype differences in *Vallisneria americana*? *Bull.Ecol. Soc. Am.* 74 (Suppl.):439.

South Florida Water Management District. 2003. Technical Documentation to Support Development of Minimum Flows and Levels for the Caloosahatchee River and Estuary, Status Update Report. May 2003.

South Florida Water Management District. 2006. South Florida Environmental Report (Draft). Chapter 12 - Caloosahatchee River, Estuary, and Southern Charlotte Harbor.

Titus, J. E. and M. S. Adams. 1979. Coexistence and the comparative light relations of the submerged macrophytes *Myriophyllum spicatum* L. and *Vallisneria americana* Michx. *Oecologia*. 40:273-286.

Wilhere, G. F. 2002. Adaptive management in habitat conservation plans. *Conserv. Biol.* 16(1): 20-29.

Xiea, Y., S. Ana, X. Yaoa, K. Xiaob, and C. Zhang. 2005. Short-time response in root morphology of *Vallisneria natans* to sediment type and water-column nutrient. Aquatic Botany 81: 85-96.