

Literature Cited

- Ceriani SA, Roth JD, Ehrhart LM, Quintana-Ascencio PF, Weishampel JF (2014a) Developing a common currency for stable isotope analyses of nesting marine turtles. *Marine Biology* 161: 2257-2268
- Ceriani SA, Roth JD, Evans DR, Weishampel JF, Ehrhart LM (2012) Inferring foraging areas of nesting loggerhead turtles using satellite telemetry and stable isotopes. *PloS one* 7: e45335 doi 10.1371/journal.pone.0045335
- Ceriani SA, Roth JD, Sasso CR, McClellan CM, James MC, Haas HL, Smolowitz RJ, Evans DR, Addison DS, Bagley DA (2014b) Modeling and mapping isotopic patterns in the Northwest Atlantic derived from loggerhead sea turtles. *Ecosphere* 5: art122-art122
- Ceriani SA, Roth JD, Tucker AD, Evans DR, Addison DS, Sasso CR, Ehrhart LM, Weishampel JF (2015a) Carry-over effects and foraging ground dynamics of a major loggerhead breeding aggregation. *Marine Biology* doi 10.1007/s00227-015-2721-x
- Ceriani SA, Weishampel JF, Ehrhart LM, KL Mansfield, Wunder MB (2015b). Modeling foraging ground dynamics at the largest loggerhead breeding aggregation in the Atlantic Ocean using species-specific isoscapes. *Proceedings from the 35th Annual Symposium on Sea Turtle Biology and Conservation*, Dalaman, Turkey, 19-24 April 2015.
- Foley AM, Schroeder BA, Hardy R, MacPherson SL, Nicholas M, Coyne MS (2013) Postnesting migratory behavior of loggerhead sea turtles *Caretta caretta* from three Florida rookeries. *Endangered Species Research* 21: 129-142 doi 10.3354/esr00512
- Hardy RF, Tucker AD, Foley AM, Schroeder BA, Giove RJ, Meylan AB. 2014. Spatiotemporal occurrence of loggerhead turtles (*Caretta caretta*) on the West Florida Shelf and apparent overlap with a commercial fishery. *Can. J. Fish. Aquat. Sci.* 71: 1924-1933
- Hobson KA (1999) Tracing Origins and Migration of Wildlife Using Stable Isotopes: A Review. *Oecologia* 120: 314-326 doi 10.2307/4222394
- Meylan PA, Meylan AB, Gray JA (2011) The ecology and migrations of sea turtles 8. Tests of the developmental habitat hypothesis. *Bull Am Mus Nat Hist* 357:1-70
- National Research Council (2010) Assessment of sea-turtle status and trends: integrating demography and abundance. Committee on the Review of Sea Turtle Population Assessment Methods (Bjorndal KA, chair). National Academies Press, Washington, D.C., USA
- National Marine Fisheries Service and US Fish and Wildlife Service (2008) Recovery plan for the Northwest Atlantic population of the loggerhead sea turtle (*Caretta caretta*). Second Revision. National Marine Fisheries Service, Silver Spring www.nmfs.gov/pr/recovery/plans.htm#turtles. Accessed 20 March 2012
- Pajuelo M, Bjorndal KA, Reich KJ, Vander Zanden HB, Hawkes LA, Bolten AB (2012) Assignment of nesting loggerhead turtles to their foraging areas in the Northwest Atlantic using stable isotopes. *Ecosphere* 3: art89 doi 10.1890/es12-00220.1
- Rubenstein DR, Hobson KA (2004) From birds to butterflies: animal movement patterns and stable isotopes. *Trends in ecology & evolution* 19: 256-263 doi 10.1016/j.tree.2004.03.017
- Schroeder BA, Foley AM, Bagley DA (2003) Nesting patterns, reproductive migrations, and adult foraging areas of loggerhead turtles. *Loggerhead sea turtles*: 114-124
- Tucker AD, MacDonald BD, Seminoff JA (2014) Foraging site fidelity and stable isotope values of loggerhead turtles tracked in the Gulf of Mexico and northwest Caribbean. *MARINE ECOLOGY PROGRESS SERIES* 502: 267-279
- Vander Zanden HB, Pfaller JB, Reich KJ, Pajuelo M, Bolten AB, Williams KL, Frick MG, Shamblin BM, Nairn CJ, Bjorndal KA (2014) Foraging areas differentially affect reproductive output and interpretation of trends in abundance of loggerhead turtles. *Marine Biology* 161: 585-598 doi 10.1007/s00227-013-2361-y

Vander Zanden HB, Tucker AD, Hart KM, Lamont MM, Fujisaki I, Addison DS, Mansfield KL, Phillips KF, Wunder MB, Bowen GJ (2015) Determining origin in a migratory marine vertebrate: a novel method to integrate stable isotopes and satellite tracking. *Ecological Applications* 25:320-335

Estimated Project Cost

Determining relative importance of foraging areas used by loggerheads nesting in Florida using stable isotope and hatchling productivity data

	Year 1	Year 2	Year 3	Year 4	Year 5	Year 6
Personnel	\$23,414	\$26,281	\$42,532	\$27,321	\$27,841	\$28,361
Fringe Benefits	\$9,602	\$10,166	\$13,868	\$10,181	\$10,189	\$10,196
Travel	\$0	\$3,360	\$8,130	\$3,360	\$3,360	\$3,360
Equipment	\$30,000	\$0	\$0	\$0	\$0	\$0
Supplies	\$5,069	\$4,339	\$155,385	\$4,339	\$4,339	\$4,339
Other	\$14,500	\$16,500	\$40,000	\$20,500	\$13,000	\$13,000
Total Direct Costs	\$82,585	\$60,646	\$259,915	\$65,701	\$58,729	\$59,256
Total Modified Total Direct Costs	\$82,585	\$60,646	\$259,915	\$65,701	\$58,729	\$59,256
Federally-negotiated indirect-cost rate	\$14,255	\$16,441	\$70,463	\$17,812	\$15,921	\$16,064
Project Totals	\$96,839	\$77,087	\$330,377	\$83,513	\$74,650	\$75,321
GRAND TOTAL	\$737,788					