


SURFICIAL SEDIMENTS OF THE
WESTERN PORTION OF THE U.S.G.S.
1:100,000 SCALE SARASOTA QUADRANGLE

BY
GUY H. MEANS, AND
RICHARD GREEN, P.G. 1776

1997



WALTER SCHMIDT
STATE GEOLOGIST AND CHIEF

The surficial sediments and soils of the western portion of the U.S.G.S. 1:100,000 scale Sarasota Quadrangle are moderately variable in their makeup. End-member lithologies of sediments in the study area include quartz sand, silt, clay, and organic material.

For the purposes of this map, surficial sediment types were determined by examining material from within the upper two meters of the land surface. Data utilized in creating this map came from a variety of sources, including the United States Department of Agriculture (U.S.D.A.) soil surveys of Manatee, Sarasota, and Charlotte Counties (Hyde and Huckle, 1983; Hyde, et al., 1991; and Henderson, 1984), hand-augered samples, surficial samples, well cuttings, and cores.

Over 50 different soil classifications were identified for the Sarasota quadrangle in the U.S.D.A. soil surveys for Manatee, Sarasota, and Charlotte Counties. For the purposes of this map, these soil types were combined into six units. These units are based on the lithologies of the constituent components of the soil types and did not take into account factors such as moisture content and plasticity. For more detailed information on soil types in the area, the reader should refer to the U.S.D.A. Soil Surveys for Manatee, Sarasota and Charlotte Counties (see Hyde and Huckle, 1983; Hyde et al., 1991; and Henderson, 1984, respectively).

The predominant constituent of the surficial sediments throughout the quadrangle is quartz sand. This is most evident in the northeastern part of the study area, where elevations are relatively high, and the undifferentiated Quaternary (Qu) sediments are thickest (see accompanying geologic map). All of the sediment types shown on this map contain at least some quartz sand. In some areas, the unit mapped as sand contains minor shell material. Although much of the area mapped as Tertiary/Quaternary undifferentiated fossiliferous sediments (TQu) on the accompanying geologic map does contain significant amounts of shell material, it was generally deeper than two meters below the land surface.

The next most common sediment type in the study area is loam. Loam, which is defined as "soil material that is 7 to 27 percent clay particles, 28 to 50 percent silt particles, and less than 52 percent sand particles" (Hyde, et al., 1991, p. 97), occurs throughout much of the map area, particularly in drainage features and basins. Occurrences of loam are most prevalent in areas where the Qu is relatively thin (see geologic map).

The remaining units depicted on this map: sand/loam, clay/sand/loam, muck, and sand/rock, are relatively minor in areal extent. Mixtures of sand and loam are most prevalent in the southern part of the study area. Clayey sediments, with admixtures of sand and loam, are present near the southern Myakka River basin. Muck, soil material which contains slowly decaying organic matter and may contain some silt and clay, is present around the Myakka River basin, the Braden River basin, and in other areas where drainage is poor. The unit mapped as sand/rock generally coincides with areas where the Arcadia Formation is within a few meters of the land surface, such as around the Braden River and Bradenton.

Selected Bibliography

Barr, G.L., 1996, Hydrogeology of the Surficial and Intermediate Aquifer Systems in Sarasota and adjacent counties, Florida: U.S. Geological Survey Water-Resources Investigations Report 96-4063, 81p.

Brewster-Wingard, G.L., Scott, T.M., Edwards, L.E., and Weedman, S.D., 1997, Reinterpretation of the peninsular Florida Oligocene: A multidisciplinary view. *Sedimentary Geology*, v.108, p.207-228.

Campbell, K.M., 1985, Geology of Sarasota County, Florida: Florida Geological Survey Open-File Report 10, 16p.

Campbell, K.M., 1993, Geologic map of Manatee County, Florida: Florida Geological Survey Open-File Map Series 50.

Campbell, K.M., and Arthur, J.D., 1993, Geologic map of Sarasota County, Florida: Florida Geological Survey Open-File Map Series 57.

Campbell, K.M., Scott, T.M., Green, R.C., and Evans, W.L. III, 1993, Sarasota County intermediate aquifer system core drilling and analysis: Florida Geological Survey Open-File Report 56, 21p.

Duerr, A.D., and Wolansky, R.M., 1986, Hydrogeology of the surficial and intermediate aquifers of central Sarasota County, Florida: U.S. Geological Survey Water-Resources Investigations Report 86-4068, 48p.

Henderson, W.G. Jr., 1984, Soil survey of Charlotte County, Florida: United States Department of Agriculture, Soil Conservation Service, 159p., 59 maps.

Hutchinson, C.B., 1992, Assessment of hydrogeologic conditions with emphasis on water quality and wastewater injection, southwest Sarasota and west Charlotte Counties, Florida: U.S. Geological Survey Water Supply Paper 2371, 74p.

Hyde, A.G., and Huckle, H., 1983, Soil survey of Manatee County, Florida: United States Department of Agriculture, Soil Conservation Service, 159p., 18 maps.

Hyde, A.G., Hurt, W., and Wettstein, C.A., 1991, Soil survey of Sarasota County, Florida: United States Department of Agriculture, Soil Conservation Service, 147p., 32 maps.

McCartan, L., and May, W., 1995, Geologic map of the Sarasota and Arcadia, Florida, 30x60 minute quadrangles: United States Geological Survey Open-File Report 95-261, 5 plates.

Petuch, E.J., 1982, Notes on the molluscan paleoecology of the Pinecrest Beds at Sarasota, Florida with the description of *Pyxulla*, a stratigraphically important new genus (Gastropoda: Melongenidae). *Proceedings of the Academy of Natural Sciences of Philadelphia*, v. 134, p. 12-30.

Petuch, E.J., 1994, Atlas of Florida fossil shells: Chicago Spectrum Press, Evanston, IL, 394 p.

Scott, T.M., 1988, The lithostratigraphy of the Hawthorn Group (Miocene) of Florida: Florida Geological Survey Bulletin 59, 148 p.

Scott, T., 1993, Coastal plains stratigraphy: The dichotomy of biostratigraphy and lithostratigraphy - A philosophical approach to an old problem: in Florida Geological Survey Special Publication 36, p. 21-26.

Scott, T.M., and Allmon, W., (eds.), 1990, Plio-Pleistocene stratigraphy and paleontology of south Florida: Southeastern Geological Society Field Trip Guidebook 31, unpaginated.

Scott, T.M., and Allmon, W., (eds.), 1993, Plio-Pleistocene stratigraphy and paleontology of southern Florida: Florida Geological Survey Special Publication 36, 194 p.


Scott, T.M., and Missimer, T., 1993, Geologic map of Charlotte County, Florida: Florida Geological Survey Open-File Map Series 59.

Scott, T.M., and Wingard, G.L., 1995, Facies, fossils, and time - A discussion of the litho- and biostratigraphic problems in the Plio-Pleistocene sediments in southern Florida: in Scott, T.M., 1995, Compiler, Stratigraphy and paleontology of the Plio-Pleistocene shell beds, southwest Florida: Southeastern Geological Society Guidebook no. 35, unpaginated.

Slattery, J.R., Coxetter, J.G., Wylan, W.J., and Moore, J.W., Jr., 1914, Sarasota Bay, Florida, Tampa Bay to Venice: U.S. Army, Corps of Engineers, House Document No., 844, 63rd Congress, 2d Session.

White, W.A., 1970, The geomorphology of the Florida peninsula: Florida Bureau of Geology Bulletin 51, 164 p.

Wingard, G.L., Weedman, S.D., Scott, T.M., Edwards, L.E., and Green, R.C., 1994, Preliminary analysis of integrated stratigraphic data from the South Venice corehole, Sarasota County, Florida: U.S. Geological Survey Open File Report 95-3, 129 p.



COOPERATIVELY FUNDED BY THE FLORIDA GEOLOGICAL SURVEY
AND THE STATEMAP COMPONENT OF THE NATIONAL GEOLOGIC
MAPPING ACT, ADMINISTERED BY THE U.S.G.S., DEPARTMENT
OF THE INTERIOR, UNDER ASSISTANCE AWARD NUMBER
1434-HQ-96-AG-01480.