


SURFICIAL SEDIMENTS OF THE EASTERN PORTION OF THE U.S.G.S.
1:100,000 SCALE SARASOTA QUADRANGLE AND THE WESTERN
PORTION OF THE U.S.G.S. 1:100,000 SCALE ARCADIA
QUADRANGLE, SOUTH-CENTRAL FLORIDA

BY
GUY H. MEANS
AND
RICHARD GREEN, P.G. # 1776
1998



WALTER SCHMIDT
STATE GEOLOGIST AND CHIEF

Surficial sediments and soils of the eastern portion of the U.S.G.S. 1:100,000 scale Sarasota and western portion of the U.S.G.S. 1:100,000 scale Arcadia quadrangles are comprised primarily of four lithologic components: quartz sand, loam, clay, and muck. Combinations of these components constitute the five lithologic units represented on this map.

Data utilized in creating this map was derived from a variety of sources, including United States Department of Agriculture Soil Surveys (Hyde & Huckle, 1983; Hyde et al., 1991; Henderson, 1984; Robbins, et al., 1984; and Cowherd, et al., 1989), field observations, cores and well cuttings, surface samples, and hand-augered samples. Sediment types shown on the map represent soil from within the upper two meters (6.6 feet) of land surface.

Between 40 and 50 different soil types are recognized, depending on the county, by the United States Department of Agriculture in published soil surveys (Hyde & Huckle, 1983; Hyde et al., 1991; Henderson, 1984; Robbins, et al., 1984; and Cowherd, et al., 1989). In these soil surveys, soil types were categorized based on factors such as slope, moisture content, texture, distribution of organic matter, acidity and other physical properties. The units which appear on this map were chosen based primarily on lithology and represent combinations of numerous soil types. Therefore, the reader is urged to consult the soil surveys for more detailed information about soils in the map area.

Quartz sand, the most commonly occurring sediment type in the map area, generally correlates to regions of higher elevation where undifferentiated Quaternary (Qu) sediments have accumulated. All of the sediment types shown on this map contain at least some quartz sand. In some areas, the unit mapped as sand may contain minor percentages of shell, phosphate sand, and heavy minerals. Although much of the area mapped as Tertiary/Quaternary undifferentiated fossiliferous sediments (TQsu) on the accompanying geologic map does contain significant amount of shell material, it is generally deeper than two meters below land surface.

Loam, defined as "soil material that contains 7 to 27 percent clay particles, 28 to 50 percent silt particles, and less than 52 percent sand size particles" (Hyde, et al., 1991, p. 97), occurs throughout much of the map area and is generally associated with drainage features and basins. Occurrences of loam are most prevalent where the Qu is relatively thin.



Clayey sediments with admixtures of sand and loam are predominately found within the Peace River and Horse Creek floodplains, with minor occurrences in other parts of the map area. This unit may correspond, in some areas, to Peace River Formation outcrops (see accompanying geologic map). This association is not always exclusive, however, since other geologic units within the quadrangle are known to contain clay.

The remaining units depicted on this map: sand/loam and muck, are relatively minor in areal extent. Mixtures of sand and loam are most prevalent in the area between the Peace River and Horse Creek in Hardee county, in the eastern portion of Hardee County, and in south-central DeSoto and north-central Charlotte Counties.

Muck, which is slowly decaying organic material containing varying amounts of sand and silt, has accumulated in large areas of the southern portion of the Peace River basin just above Charlotte Harbor. Several areas in southeastern DeSoto County including Tiger Bay, Sheep Pen Marsh, and Long Island Marsh also contain muck. These are generally swampy, poorly drained areas which allow organic material to accumulate.

Selected Bibliography

- Arthur, J.D., 1993, Geologic map of DeSoto County, Florida: Florida Geological Survey Open-File Map Series 58.
- Brewster-Wingard, G.L., Scott, T.M., Edwards, L.E., and Weedman, S. D., 1987, Reinterpretation of the peninsular Florida Oligocene: A multidisciplinary view: Sedimentary Geology, v.108, p. 207-228.
- Campbell, K.M., 1985, Geology of Sarasota County, Florida: Florida Geological Survey Open-File Report 10, 15p., 6 fig.
- Campbell, K.M., 1985, Geology of DeSoto County, Florida: Florida Geological Survey Open-File Report 11, 13p., 4 fig.
- Campbell, K.M., 1993, Geologic map of Manatee County, Florida: Florida Geological Survey Open-File Map Series 50.
- Campbell, K.M., and Arthur, J.D., 1993, Geologic map of Sarasota County, Florida: Florida Geological Survey Open-File Map Series 57.
- Cowherd, D.W., Henderson, W.G., Sheehan, E.J. and Ploetz, S.T., 1989, Soil survey of DeSoto County, Florida: United States Department of Agriculture, Soil Conservation Service, 170p., 15 maps.
- Green, R., Campbell, K., Arthur, J., and Scott, T., 1997, Geologic map of the western portion of the U.S.G.S. 1:100,000 scale Sarasota quadrangle: Florida Geological Survey Open-File Map Series 86, 8 plates.
- Henderson, W.G. Jr., 1984, Soil survey of Charlotte County, Florida: United States Department of Agriculture, Soil Conservation Service, 159p., 18 maps.
- Hyde, A.G., and Huckle, H.F., 1983, Soil survey of Manatee County, Florida: United States Department of Agriculture, Soil Conservation Service, 159p., 17 maps.
- Hyde, A.G., Hurt, W., and Wettstein, C.A., 1991, Soil survey of Sarasota County, Florida: United States Department of Agriculture, Soil Conservation Service, 147p., 32 maps.
- Lowell, B.R., Tihonsky, A.B. and Kindinger, J.L., 1998, Assessment of the hydraulic connection between groundwater and the Peace River, west-central Florida: United States Geological Survey Water Resources Investigations Report 97-4211, 96p.
- McCartan, L., and Moy, W., 1995, Geologic map of the Sarasota and Arcadia, Florida, 30x60 minute quadrangles: United States Geological Survey Open-File Report 95-261, 5 plates.
- Petuch, E.J., 1982, Notes on the molluscan paleoecology of the Pliocene Beds at Sarasota, Florida with the description of *Eryxella*, a stratigraphically important new genus (Gastropoda: Melanogastropoda): Proceedings of the Academy of Natural Sciences of Philadelphia, v. 134, p.12-30.
- Robbins, J.M., Ford, R.D., Werner, J.T., and Cowherd, W.D., 1984, Soil survey of Hardee County, Florida: United States Department of Agriculture, Soil Conservation Service, 139p., 49 maps.
- Scott, T.M., 1988, The lithostratigraphy of the Hawthorn Group (Miocene) of Florida, Florida Geological Survey Bulletin 59, 148p.
- Scott, T.M., 1993, Coastal plains stratigraphy: The dichotomy of biostratigraphy and lithostratigraphy - A philosophical approach to an old problem: In Florida Geological Survey Special Publication 36, p. 21-26.
- Scott, T.M., 1998, (In preparation), Geomorphologic map of Florida: Florida Geological Survey.
- Scott, T.M., (editor and compiler), Campbell, K.M., Rupert, F.R., Arthur, J., Duncan, J., Missimer, T., Lloyd, J., and Yon, J.W., 1998 (In preparation), Geologic map of Florida: Florida Geological Survey.
- Scott, T.M., Allmon, W., 1990, Plio-Pleistocene stratigraphy and paleontology of south Florida: Southeastern Geological Society Field Trip Guidebook 31, unpaginated.
- Scott, T.M., and Allmon, W., (eds.), 1993, Plio-Pleistocene stratigraphy and paleontology of south Florida: Florida Geological Survey Special Publication 36, 194p.
- Scott, T.M., and Campbell, K.M., 1993, Geologic map of Hardee County, Florida: Florida Geological Survey Open-File Map Series 51.
- Scott, T.M., and Missimer, T., 1993, Geologic map of Charlotte County, Florida: Florida Geological Survey Open-File Map Series 59.
- Scott, T.M., and Wingard, G.L., 1995, Facies, fossils, and time - A discussion of the litho- and biostratigraphic problems in the Plio-Pleistocene sediments of southern Florida: In Scott, T.M., 1995, Compiler, Stratigraphy and paleontology of the Plio-Pleistocene shell beds, southwest Florida: Southeastern Geological Society Guidebook no. 35, unpaginated.



COOPERATIVELY FUNDED BY THE FLORIDA GEOLOGICAL SURVEY AND THE STATEMAP COMPONENT OF THE NATIONAL GEOLOGIC MAPPING ACT, ADMINISTERED BY THE U.S.G.S., DEPARTMENT OF THE INTERIOR, UNDER ASSISTANCE AWARD NUMBER 1434-HQ-97-AG-01808

